



Brunei International Medical Journal

OFFICIAL PUBLICATION OF
THE MINISTRY OF HEALTH
AND
UNIVERSITI BRUNEI DARUSSALAM

Volume 18

15 April 2022 (13 Ramadan 1443H)

SMARTPHONE USAGE AND PATTERNS TOWARDS SELF-REPORTED SYMPTOMS DURING EARLY 2019 CORONAVIRUS PANDEMIC LOCKDOWN AMONG MEDICAL STUDENTS IN MALAYSIA.

Megat KAMARUDDIN PSN, Mohammed NAWI A.

Department of Community Health, Faculty of Medicine, Universiti Kebangsaan Malaysia, Bandar Tun Razak, Kuala Lumpur, Malaysia.

ABSTRACT

Introduction: Smartphone ubiquity has become profound during coronavirus disease (COVID-19). Prior research suggests smartphone usage and patterns correlated with self-reported symptoms. This study aimed to evaluate smartphone usage and associated patterns of self-reported symptoms among medical students in Malaysia during COVID-19 pandemic lockdown. **Materials and Methods:** This was a cross-sectional study of 252 medical students who were given questionnaires regarding sociodemographic, socio-economic characteristics, smartphone usage and patterns of self-reported symptoms. Total severity scores of self-reported symptoms were calculated. Association between socio-demographic, socio-economic and academic performance status with smartphone usage and patterns were examined using simple and multiple linear regression analysis. **Results:** Mean Smartphone Addiction Scale–Malay score for smartphone usage was 101.43 (SD = 25.15). Smartphone patterns revealed 7 years of smartphone ownership, 32 minutes per day on calls and 4 hours per day on others (for entertainment (50%) and non-entertainment (50%) purposes. They had one extra device and used their smartphone 15 minutes after waking up in the morning. After controlling other factors, father's employment status (adjusted (Adj.) $\beta=7.431$, 95% CI: 3.069, 11.793) and smartphone usage for entertainment purposes (Adj. $\beta = 4.211$, 95% CI: 0.460, 7.962) were only two significant predictors for self-reported symptoms. **Conclusion:** Father's employment status and smartphone usage for entertainment purposes were associated with increase frequency of self-reported symptoms among medical students during the early lockdown of the COVID-19 pandemic in Malaysia. Parents' participation from early childhood and public campaigns to avoid smartphone addiction are essential.

Keywords: COVID-19, Medical student(s), Pattern(s), Symptoms, Smartphone.

Brunei Int Med J. 2022;18:63-74

Brunei International Medical Journal (BIMJ)

Official Publication of The Ministry of Health and Universiti Brunei Darussalam

EDITORIAL BOARD

Editor-in-Chief	Ketan PANDE
Sub-Editors	Vui Heng CHONG William Chee Fui CHONG
Editorial Board Members	Muhd Syafiq ABDULLAH Alice Moi Ling YONG Ahmad Yazid ABDUL WAHAB Jackson Chee Seng TAN Pemasiri Upali TELISINGHE Pengiran Khairol Asmee PENGIRAN SABTU Dayangku Siti Nur Ashikin PENGIRAN TENGAH

INTERNATIONAL EDITORIAL BOARD MEMBERS

Lawrence HO Khek Yu (Singapore)	Chuen Neng LEE (Singapore)
Wilfred PEH (Singapore)	Emily Felicia Jan Ee SHEN (Singapore)
Surinderpal S BIRRING (United Kingdom)	Leslie GOH (United Kingdom)
John YAP (United Kingdom)	Ian BICKLE (United Kingdom)
Nazar LUQMAN (Australia)	Christopher HAYWARD (Australia)
Jose F LAPENA (Philippines)	

Advisor

Wilfred PEH (Singapore)

Past Editors-in-Chief

Nagamuttu RAVINDRANATHAN
Kenneth Yuh Yen KOK
Chong Vui Heng
William Chong Chee Fui

Proof reader

John WOLSTENHOLME (CfBT Brunei Darussalam)

Aim and Scope of Brunei International Medical Journal

The Brunei International Medical Journal (BIMJ) is a six monthly peer reviewed official publication of the Ministry of Health under the auspices of the Clinical Research Unit, Ministry of Health, Brunei Darussalam.

The BIMJ publishes articles ranging from original research papers, review articles, medical practice papers, special reports, audits, case reports, images of interest, education and technical/innovation papers, editorials, commentaries and letters to the Editor. Topics of interest include all subjects that relate to clinical practice and research in all branches of medicine, basic and clinical including topics related to allied health care fields. The BIMJ welcomes manuscripts from contributors, but usually solicits reviews articles and special reports. Proposals for review papers can be sent to the Managing Editor directly. Please refer to the contact information of the Editorial Office.

Instruction to authors Manuscript submissions

All manuscripts should be sent to the Managing Editor, BIMJ, Ministry of Health, Brunei Darussalam; e-mail: editor-in-chief@bimjonline.com. Subsequent correspondence between the BIMJ and authors will, as far as possible via should be conducted via email quoting the reference number.

Conditions

Submission of an article for consideration for publication implies the transfer of the copyright from the authors to the BIMJ upon acceptance. The final decision of acceptance rests with the Editor-in-Chief. All accepted papers become the permanent property of the BIMJ and may not be published elsewhere without written permission from the BIMJ.

Ethics

Ethical considerations will be taken into account in the assessment of papers that have experimental investigations of human or animal subjects. Authors should state clearly in the Materials and Methods section of the manuscript that institutional review board has approved the project. Those investigators without such review boards should ensure that the principles outlined in the Declaration of Helsinki have been followed.

Manuscript categories

Original articles

These include controlled trials, interventional studies, studies of screening and diagnostic tests, outcome studies, cost-effectiveness analyses, and large-scale epidemiological studies. Manuscript should include the following; introduction, materials and methods, results and conclusion. The objective should be stated clearly in the introduction. The text should not exceed 2500 words and references not more than 30.

Review articles

These are, in general, invited papers, but unsolicited reviews, if of good quality, may be considered. Reviews are systematic critical assessments of

literature and data sources pertaining to clinical topics, emphasising factors such as cause, diagnosis, prognosis, therapy, or prevention. Reviews should be made relevant to our local setting and preferably supported by local data. The text should not exceed 3000 words and references not more than 40.

Special Reports

This section usually consist of invited reports that have significant impact on healthcare practice and usually cover disease outbreaks, management guidelines or policy statement paper.

Audits

Audits of relevant topics generally follow the same format as original article and the text should not exceed 1,500 words and references not more than 20.

Case reports

Case reports should highlight interesting rare cases or provide good learning points. The text should not exceed 1000 words; the number of tables, figures, or both should not be more than two, and references should not be more than 15.

Education section

This section includes papers (i.e. how to interpret ECG or chest radiography) with particular aim of broadening knowledge or serve as revision materials. Papers will usually be invited but well written paper on relevant topics may be accepted. The text should not exceed 1500 words and should include not more than 15 figures illustration and references

three relevant references should be included. Only images of high quality (at least 300dpi) will be acceptable.

Technical innovations

This section include papers looking at novel or new techniques that have been developed or introduced to the local setting. The text should not exceed 1000 words and should include not more than 10 figures illustration and references should not be more than 10.

Letters to the Editor

Letters discussing a recent article published in the BIMJ are welcome and should be sent to the Editorial Office by e-mail. The text should not exceed 250 words; have no more than one figure or table, and five references.

Criteria for manuscripts

Manuscripts submitted to the BIMJ should meet the following criteria: the content is original; the writing is clear; the study methods are appropriate; the data are valid; the conclusions are reasonable and supported by the data; the information is important; and the topic has general medical interest. Manuscripts will be accepted only if both their contents and style meet the standards required by the BIMJ.

Authorship information

Designate one corresponding author and provide a complete address, telephone and fax numbers, and e-mail address. The number of authors of each paper should not be more than twelve; a greater number requires justification. Authors may add a publishable footnote explaining order of authorship.

Group authorship

If authorship is attributed to a group (either solely or in addition to one or more individual authors), all members of the group must meet the full criteria and requirements for authorship described in the following paragraphs. One or more authors may take responsibility 'for' a group, in which case the other group members are not authors, but may be listed in an acknowledgement.

Authorship requirement

DISCLAIMER

All articles published, including editorials and letters, represent the opinion of the contributors and do not reflect the official view or policy of the Clinical Research Unit, the Ministry of Health or the institutions with which the contributors are affiliated to unless this is clearly stated. The appearance of advertisement does not necessarily constitute endorsement by the Clinical Research Unit or Ministry of Health, Brunei Darussalam. Furthermore, the publisher cannot accept responsibility for the correctness or accuracy of the advertisers' text and/or claim or any opinion expressed.

sign, and the analysis and interpretation of the data (where applicable); to have made substantial contributions to the writing or revision of the manuscript; and to have reviewed the final version of the submitted manuscript and approved it for publication. Authors will be asked to certify that their contribution represents valid work and that neither the manuscript nor one with substantially similar content under their authorship has been published or is being considered for publication elsewhere, except as described in an attachment. If requested, authors shall provide the data on which the manuscript is based for examination by the editors or their assignees.

Financial disclosure or conflict of interest

Any affiliation with or involvement in any organisation or entity with a direct financial interest in the subject matter or materials discussed in the manuscript should be disclosed in an attachment. Any financial or material support should be identified in the manuscript.

Copyright transfer

In consideration of the action of the BIMJ in reviewing and editing a submission, the author/s will transfer, assign, or otherwise convey all copyright ownership to the Clinical Research Unit, RIPAS Hospital, Ministry of Health in the event that such work is published by the BIMJ.

Acknowledgements

Only persons who have made substantial contributions but who do not fulfill the authorship criteria should be acknowledged.

Accepted manuscripts

Authors will be informed of acceptances and accepted manuscripts will be sent for copyediting. During copyediting, there may be some changes made to accommodate the style of journal format. Attempts will be made to ensure that the overall meaning of the texts are not altered. Authors will be informed by email of the estimated time of publication. Authors may be requested to provide raw data, especially those presented in graph such as bar charts or figures so that presentations can be constructed following the format and style of the journal. Proofs will be sent to authors to check for any mistakes made

SMARTPHONE USAGE AND PATTERNS TOWARDS SELF-REPORTED SYMPTOMS DURING EARLY 2019 CORONAVIRUS PANDEMIC LOCKDOWN AMONG MEDICAL STUDENTS IN MALAYSIA.

Megat KAMARUDDIN PSN, Mohammed NAWI A.

Department of Community Health, Faculty of Medicine, Universiti Kebangsaan Malaysia, Bandar Tun Razak, Kuala Lumpur, Malaysia.

ABSTRACT

Introduction: Smartphone ubiquity has become profound during coronavirus disease (COVID-19). Prior research suggests smartphone usage and patterns correlated with self-reported symptoms. This study aimed to evaluate smartphone usage and associated patterns of self-reported symptoms among medical students in Malaysia during COVID-19 pandemic lockdown. **Materials and Methods:** This was a cross-sectional study of 252 medical students who were given questionnaires regarding sociodemographic, socioeconomic characteristics, smartphone usage and patterns of self-reported symptoms. Total severity scores of self-reported symptoms were calculated. Association between socio-demographic, socio-economic and academic performance status with smartphone usage and patterns were examined using simple and multiple linear regression analysis. **Results:** Mean Smartphone Addiction Scale-Malay score for smartphone usage was 101.43 (SD = 25.15). Smartphone patterns revealed 7 years of smartphone ownership, 32 minutes per day on calls and 4 hours per day on others (for entertainment (50%) and non-entertainment (50%) purposes. They had one extra device and used their smartphone 15 minutes after waking up in the morning. After controlling other factors, father's employment status (adjusted (Adj.) $\beta=7.431$, 95% CI: 3.069, 11.793) and smartphone usage for entertainment purposes (Adj. $\beta = 4.211$, 95% CI: 0.460, 7.962) were only two significant predictors for self-reported symptoms. **Conclusion:** Father's employment status and smartphone usage for entertainment purposes were associated with increase frequency of self-reported symptoms among medical students during the early lockdown of the COVID-19 pandemic in Malaysia. Parents' participation from early childhood and public campaigns to avoid smartphone addiction are essential.

Keywords: COVID-19, Medical student(s), Pattern(s), Symptoms, Smartphone.

INTRODUCTION

There has been an upsurge in research into smartphone use among young people, but

little on long-term monitoring of their potential health effects.¹ Non-ionising radiation penetration into their brains may consequently change the permeability of the blood-brain barrier, which, young individuals are believed to be more prone to symptoms of smartphone usage.² A powerful electromagnetic field (EMF) may induce illnesses such as insomnia, headaches, and allergy-like symptoms, known as

Corresponding author: Azmawati Mohammed Nawi, (MD, Mcomm. Med, PhD), Department of Community Health, Faculty of Medicine, Universiti Kebangsaan Malaysia, Jalan Yaacob Latif, Bandar Tun Razak, 56000 Kuala Lumpur, Malaysia.
Tel: +603-91458780. Fax: +603-91456670.
Email: azmawati@ppukm.ukm.edu.my

Electrical Hypersensitivity (EHS), although no definite causal link between EMF exposure and symptoms has been shown.³ In two studies conducted on junior high school and university students respectively, frequent smartphone usage has been reported to be associated with headaches, memory loss, insomnia, and other mental disorders.^{4,5} Students' coping techniques and priorities, such as coursework, may go unrecognised masking the health symptoms.²

On March 18th 2020, Malaysian government imposed measures to limit viral transmission and decrease contact with affected patients of coronavirus disease 2019 (COVID-19).⁶ The global pandemic has affected the operational activities of educational institutions, which require pupils to carry out online studies at home and may inadvertently increase their access and exposure to mobile technology. Hence self-reporting of symptoms such as headaches, insomnia, memory loss etc may be expected to increase during the lockdown period.

The aim of this study was to investigate the frequency and patterns of smartphone usage among medical students at Universiti Kebangsaan Malaysia (UKM) and the association of self-reported symptoms during the COVID-19 pandemic lockdown. We hypothesised a higher smartphone usage and an associated increase in self-reported symptoms during the period of COVID-19 pandemic lockdown.

MATERIALS AND METHODS

From 16th March to 14th April 2020, a cross-sectional study was performed among UKM undergraduate Medical Degree students with smartphones, utilising proportionate stratified random sampling, in the setting of COVID-19 lockdown.⁶ The UKM Ethics Committee granted ethical approval (FF-2020-105). All partici-

pants gave written consent for participation and were divided into groups based on their clinical year and exposure: preclinical (first and second years) and clinical (third, fourth and fifth years), by randomly generated students' matriculation numbers. Medical students with psychiatric disorders, cancer, thyroid disease, anaemia, explicit drug use, pregnant or hospitalised for any reason were excluded.

Utilising Kish equation, the sample size derived was 252, with 10% for missing data or drop-out, based on self-reported physical symptoms value of 'Pain in the neck muscle, 58.5%, n = 134'.^{7,8} This study's sample size was determined using 80% power and a 0.05 of type I error. Where precision was needed, the 95% CI and 5% precision were used.

Study tools

Utilising a web-based Google sheet form to record the responses, the first part included data on sociodemographic and socioeconomic factors, such as age, ethnic, parents' income, occupation, and educational levels, as well as the student's funding status and academic achievement. The second segment focused on smartphone usage patterns, including the amount of smartphone use on a regular day, the length of smartphone ownership, the time between wake up and smartphone initial use, the number of additional gadgets owned, and the most important smartphone function.

Smartphone usage details were requested in the third section. The Smartphone Addiction Scale-Malay (SAS-M) was utilised, which previously created, translated, and validated among Malaysian medical students, with Cronbach's alpha of 0.94.⁹ The SAS-M features six subscales (cyberspace-oriented connection, everyday life disturbance, primacy, overuse, etc.) of 33 items. Cronbach's alpha was 0.91 among 30 participants, while the corresponding coefficients for the six vari-

ables were 0.78, 0.76, 0.86, 0.67, 0.83, and 0.85. Each item is scored on 1–6 scale (1 = strongly disagree; 6 = strongly agree), with scoring of 33–198, which higher values suggesting likely of smartphone addiction or use.⁹

The fourth segment dealt with self-reported symptoms. Each student declared the ten symptoms encountered and assessed their severity on a scale of 0–10, with 0 representing none and 10 representing often everyday. Using the Total Score of Severity (TSS) equation¹¹:

- TSS per person = Σ (self-reported symptom $\times$ symptom's severity factor)

TSS is the sum of ten self-reported symptoms experienced. TSS ratings range from 0 to 100, with higher values indicating more severe symptoms.¹⁰

Statistical analysis

Continuous variables were summarised using mean and standard deviation (SD) and categorical variables were summarised using frequencies and percentages (%). Using SPSS version 25 (Statistical Package for Social Sciences), data on smartphone usage and patterns were analysed using Student's t-test, Pearson's chi square, and Pearson's correlation. Using simple and multivariate linear regression (MLR), we found the main component contributing to the association between smartphone usage and patterns with self-reported symptoms. The significance level for all tests was 0.05. All tests were two-sided, and $p < 0.05$ was deemed significant.

RESULTS

The characteristics of the participants (Table I) reported that the mean age (SD) was 23.09 (1.84) years; 75.4% were females, and 59.9% were of Malay origin. The partici-

Table I: Participant's characteristics, smartphone usage and pattern.

Participant's Characteristics	n (%) N = 252	(Mean (SD))
Age (years)		23.09 (1.84)
Gender		
Male	62 (24.6)	
Female	190 (75.4)	
Ethnicity		
Malay	151 (59.9)	
Chinese	47 (18.7)	
Indian	42 (16.7)	
^a Others	12 (4.8)	
Parents' income (RM)		6641.57 (7067.56)
Father's occupational status		
Employed	190 (75.4)	
Unemployed	62 (24.6)	
Mother's occupation status		
Employed	136 (54.0)	
Unemployed	116 (46.0)	
Father's education level		
Nil	2 (0.8)	
Primary	18 (7.1)	
Secondary	77 (30.6)	
Diploma / Certificate	54 (21.4)	
Tertiary	101 (40.1)	
Mother's education level		
Nil	4 (1.6)	
Primary	15 (6.0)	
Secondary	97 (38.5)	
Diploma / Certificate	53 (21.0)	
Tertiary	83 (32.9)	
Funding status		
Yes	49 (19.4)	
No	203 (80.6)	
Academic achievement (CGPA)		3.2 (0.37)
Smartphone usage and patterns		
Smartphone usage		
SAS-M score		101.43 (25.15)
Smartphone patterns		
Most personal smartphone feature		
Entertainment	126 (50.0)	
Non-entertainment	126 (50.0)	
Time spent making or receiving phone calls per day (minutes)		32.17 (49.97)
Non-call related smartphone usage per day (hours)		4.26 (4.29)
Ownership of smartphone (years)		7.23 (2.77)
Time from waking to first smartphone usage in the morning (minutes)		14.68 (34.92)
Number of additional devices owned		1.38 (0.68)

Abbreviations: SD standard deviation, RM Ringgit Malaysia
^aOthers, n (%): Bumiputera 3 (1.2), Kadazan 2 (0.8), Bajau 1 (0.4), Bidayuh 1 (0.4), Bugis 1 (0.4), Dusun 1 (0.4), Punjabi 1 (0.4), Siamese 1 (0.4), Sino-Kadazan 1 (0.4)

pants' parents earned an average (SD) of RM6641.57 (RM7067.56) a month in gross monthly earnings; 40.1% of the participants' fathers had achieved tertiary education, while 38.5% of the participants' mothers had completed secondary school. The majority of respondents had working parents: 75.4% of fathers and 54% of mothers. Approximately 80.6% of the participants were not obtaining any aid or loan for their present studies, and the previous semester's cumulative grade point average CGPA (SD) was 3.2 (0.37).

The participants' average period of smartphone ownership was 7 years with an average (SD) SAS-M score for smartphone usage (Table I) of 101.43 (25.15) and utilisation of smartphones were equally divided for entertainment and non-entertainment purposes (50%). They made or received calls on average for around 32 minutes per day, and that they spent approximately 4 hours per day on smartphone usage other than calls. They had one additional gadget in addition to their smartphone, and they used their smartphone within 15 minutes of waking up in the morning.

In terms of self-reporting of symptoms, majority of the medical students scored

zero (range 24.6-81.3%) and did not report any symptoms. The three most common self-reported symptoms were insomnia (75.4%), neck pain (67.9%) and headache (63.9%), although in the majority, these were either mild or moderate in rating scale (Table II:). Less than a third complaint of moderate symptoms scale (4-6), of which majority were reporting of insomnia (Table II: 29.3%). Eleven percent or less of participants reported moderately high symptom score (7-8) and very few (range 0.4 to 4.4%) reported high symptom score (Scale 9-10). Overall, the average TSS (SD) for all reported symptoms was low at 17.84 (15.58) out of a 100 (Table II).

There was a significant association between being of Malay ethnicity, having employed and well-educated parents, participant's academic achievement with increased smartphone usage (Table III) and to various smartphone usage patterns (Table IV).

After adjusting for other variables, multivariable analysis revealed that father with employment status and using smartphone for entertainment function were the significant predictors of higher TSS (Table V). Medical students whose father

Table II: Self-reported symptoms among medical students.

Self – reported symptoms	Rating Scale n (%)					Mean (SD)
	0	1-3 (Mild)	4-6 (Moderate)	7-8 (Moderately high)	9-10 (High)	
Headache	91 (36.1)	106 (42.1)	42 (16.76)	12 (4.8)	1 (0.4)	
Dizziness	111 (44.0)	93 (36.9)	38 (15.1)	9 (3.6)	1 (0.4)	
Neck pain	81 (32.1)	94 (37.3)	49 (19.4)	22 (8.8)	6 (2.4)	
Hearing problem	161 (63.9)	54 (21.4)	30 (12.0)	5 (2.0)	2 (0.8)	
Vision problem	99 (39.3)	81 (32.1)	47 (18.6)	21 (8.4)	4 (1.6)	
Insomnia	62 (24.6)	77 (30.5)	74 (29.3)	28 (11.1)	11 (4.4)	
Palpitation	172 (68.3)	53 (21.0)	17 (6.8)	8 (3.2)	2 (0.8)	
General exhaustion	109 (43.3)	75 (29.7)	44 (17.5)	17 (6.8)	7 (2.8)	
Shortness of breath	205 (81.3)	40 (15.9)	5 (2.0)	0	2 (0.8)	
Temporary memory loss	156 (61.9)	65 (25.7)	24 (9.5)	4 (1.6)	3 (1.2)	
Total Score of Severity (TSS)						17.84 (15.58)

Abbreviations: SD standard deviation, SAS-M Smartphone Addiction Scale-Malay version, TSS Total Score of Severity

Table III: Smartphone usage across sample characteristics.

Variables	Smartphone usage Mean (SD)	(SAS-M score) t / r	p value
Sociodemography			
Age (years)		-0.015 ^a	0.814
Gender			
Male	100.9 (20.86)	.213 ^b	0.832
Female	101.6 (26.45)		
Ethnicity			
Malay	115.43 (16.56)	14.75 ^b	<0.001*
Non-Malay	80.50 (20.92)		
Socioeconomy			
Parents' income (RM)	-	-0.003 ^a	0.960
Father's Occupational Status			
Employed	107.53 (22.53)	7.430 ^b	<0.001*
Unemployed	82.74 (23.63)		
Mother's Occupational Status			
Employed	119.82 (16.29)	20.601 ^b	<0.001*
Unemployed	79.87 (14.15)		
Father's education level			
High	106.48 (29.04)	4.81 ^b	<0.001*
Low	93.36 (13.93)		
Mother's education level			
High	105.48 (32.62)	3.001 ^b	0.003*
Low	96.68 (9.46)		
Funding status			
Yes	100.81 (25.74)	0.791 ^b	0.430
No	103.98 (22.61)		
Academic achievement	-	0.987 ^a	<0.001*

^ar = Pearson's correlation coefficient; ^bt = Student's t-test value

Abbreviations: SD standard deviation, RM Ringgit Malaysia

*p value = statistically significant at <0.05

were employed at the time of study were seven times more likely to self-report of symptoms, (Table V: adjusted $\beta = 7.431$, 95% CI: 3.069, 11.793), and using their smartphone for entertainment function, was associated with a four times increase in self-reported symptoms (Table V: Adj. $\beta = 4.211$, 95% CI: 0.460, 7.962).

DISCUSSION

The SAS-M score (SD) of 101.43 (25.15) indicated the participants' smartphone usage, similar with those of previous local research, which showed marks of >98⁵ and 102.52.⁴ The results of other international studies are consistent with ours since they used the same score in modified Spanish, Arabic, and

Turkish, which higher score indicates a greater risk of smartphone use or addiction.^{11, 12} During the COVID-19 pandemic, medical students are expected to rely heavily on technology for learning, since all Malaysian Higher Learning Institutions have transitioned to synchronous or asynchronous online learning.¹³

The average period of smartphone ownership in this survey, i.e. 7 years, is close to an Indian study that showed smartphone ownership among the majority of students was 1–8 years, since smartphones are now a necessity.¹⁴ Our results that participants made or received calls for around 32 minutes per day and spent approximately 4 hours per day on usage other than calls are similar with

Table IV: Smartphone patterns across sample characteristics.

Variables	V1		V2		V3		V4		V5		V6 ₁ /V6 ₂	
	Mean (SD)	t/r (p value)	Mean (SD)	t/r (p value)	Mean (SD)	t/r (p value)	Mean (SD)	t/r (p value)	Mean (SD)	t/r (p value)	Mean (SD) ₁ /Mean (SD) ₂	t/r (p value)
Age (Years)	-	-0.008 ^a (0.903)	-	0.029 ^a (0.649)	-	0.010 ^a (0.872)	-	0.024 ^a (0.703)	-	0.005 ^a (0.932)	23.17(2.06) / 23.0(1.59)	0.685 ^b (0.494)
Gender												
Male	28.34 (43.12)	0.694 ^b (0.488)	3.81 (2.26)	0.951 ^b (0.343)	7.76 (2.57)	1.747 ^b (0.082)	12.34 (38.47)	0.607 ^b (0.545)	1.40 (0.66)	0.348 ^b (0.728)	-	2.139 ^c (0.144)
Female	33.42 (52.05)		4.41 (4.77)		7.05 (2.82)		15.44 (33.75)		1.37 (0.69)			
Ethnicity												
Malay	39.3 (58.81)	3.163 ^b (0.002)*	5.35 (2.81)	5.157 ^b (<0.001)	7.26 (2.75)	0.224 ^b (0.823)	14.9 (36.19)	0.122 ^b (0.903)	1.40 (0.67)	0.766 ^b (0.444)	-	0.413 ^c (0.520)
Non-Malay	21.5 (29.80)		2.64 (5.47)	*	7.18 (2.81)		14.35 (33.09)		1.34 (0.71)			
Parents' income (RM)	-	0.032 ^a (0.617)	-	-0.009 ^a (0.888)	-	-0.044 ^a (0.492)	-	0.012 ^a (0.849)	-	0.047 ^a (0.454)	6601.98 (7843.97) / 6681.15 (6226.52)	0.089 ^b (0.929)
Father's occupational status												
Employed	33.41 (55.10)	0.691 ^b (0.490)	4.93 (4.68)	6.868 ^b (<0.001)	7.32 (2.79)	0.898 ^b (0.370)	13.56 (32.61)	0.894 ^b (0.372)	1.38 (0.70)	0.080 ^b (0.937)	-	0.086 ^c (0.770)
Unemployed	28.35 (29.12)		2.23 (1.56)	*	6.95 (2.73)		18.12 (41.29)		1.37 (0.63)			
Mother's occupational status												
Employed	36.75 (60.66)	1.652 ^b (0.100)	6.20 (5.03)	9.597 ^b (<0.001)	7.27 (2.74)	0.284 ^b (0.777)	14.19 (37.87)	0.239 ^b (0.811)	1.37 (0.67)	0.234 ^b (0.815)	-	0.000 ^c (1.000)
Unemployed	26.79 (32.80)		1.99 (0.91)	*	7.17 (2.82)		15.25 (31.24)		1.39 (0.71)			
Father's education level												
High	33.86 (60.44)	0.798 ^b (0.426)	5.14 (5.23)	5.259 ^b (<0.001)	7.26 (2.55)	0.277 ^b (0.782)	14.62 (41.98)	0.034 ^b (0.973)	1.26 (0.60)	3.196 ^b (0.002)*	-	0.017 ^c (0.897)
Low	29.46 (25.62)		2.86 (1.03)	*	7.16 (3.11)		14.77 (18.97)		1.56 (0.76)			
Mother's education level												
High	32.08 (59.92)	0.029 ^b (0.977)	5.26 (5.62)	4.469 ^b (<0.001)	7.01 (2.71)	1.313 ^b (0.190)	13.34 (38.40)	0.660 ^b (0.510)	1.32 (0.580)	1.498 ^b (0.136)	-	0.575 ^c (0.448)
Low	32.27 (35.15)		3.09 (0.68)	*	7.47 (2.83)		16.25 (30.41)		1.45 (0.78)			
Funding status												
Yes	30.61 (50.94)	1.009 ^b (0.314)	4.30 (4.62)	0.317 ^b (0.752)	7.30 (2.81)	0.808 ^b (0.420)	15.23 (38.11)	0.507 ^b (0.613)	1.38 (0.68)	0.342 ^b (0.732)	-	2.052 ^c (0.152)
No	38.63 (45.65)		4.09 (2.52)		6.94 (2.61)		12.41 (16.04)		1.35 (0.72)			
Academic achievement	-	0.139 ^a (0.027)*	-	0.728 ^a (<0.001)	-	0.052 ^a (0.413)	-	-0.049 ^a (0.440)	-	-0.078 ^a (0.219)	3.190 (0.35) / 3.178 (0.39)	0.257 ^b (0.797)

^ar = Pearson's correlation coefficient; ^bt = Student's t-test value; ^cχ² = Pearson's Chi-Square value; Abbreviations: SD standard deviation, RM Ringgit Malaysia
*p value = statistically significant at <0.05. **Note:** V1: Time spent making or receiving phone calls on a normal day (minutes); V2: Typical day's smartphone usage (excluding calls) (hours); V3: Time spent using a smartphone (years); V4: Time to first smartphone usage in the morning (minutes); V5: Number of additional devices owned (devices); V6₁: Most personal smartphone feature (Entertainment); V6₂: Most personal smartphone feature (Non-entertainment).

Table V: Factors associated with Total Score of Severity among medical students.

Variables	Mean (SD)	SLR ^a			MLR ^b		
		β ^c	(95% CI)	p value	Adj. β ^d	(95% CI)	p value
Sociodemography							
Age (years)	-	-0.89	(-1.95, 0.16)	0.10	-	-	-
Gender							
Male	16.58 (16.19)	-1.67	(6.17, 2.82)	0.46	-	-	-
Female	18.25 (15.40)						
Ethnicity							
Malay	19.41 (16.32)	3.92	(0.008,7.84)	0.05	-	-	-
Non-Malay	15.5 (14.17)						
Parents' income (RM)		7.83 x 10 ⁻⁵	(-0.0002,0.0004)	0.56	-	-	-
Father's Occupational Status							
Employed	19.61 (16.49)	7.19	(2.78, 11.60)	0.001*	7.43	(3.07,11.79)	0.001*
Unemployed	12.42 (10.85)						
Mother's Occupational Status							
Employed	19.19 (16.57)	2.93	(-0.94, 6.80)	0.14	-	-	-
Unemployed	16.26 (14.25)						
Father's education level							
High	17.55 (15.17)	-0.74	(-4.73, 3.24)	0.71	-	-	-
Low	18.30 (16.29)						
Mother's education level							
High	19.01 (15.51)	2.53	(-1.34, 6.41)	0.20	-	-	-
Low	16.47 (15.63)						
Funding status							
Yes	17.71 (15.59)	-0.65	(-5.55, 4.24)	0.79	-	-	-
No	18.37 (15.72)						
Academic achievement	4.49	(0.77,9.76)	0.09	-	-	-	
Smartphone usage characteristics							
SAS-M score		0.06	(-0.02, 0.13)	0.14	-	-	-
Smartphone patterns characteristics							
Time spent making or receiving phone calls per day (minutes)		0.030	(-0.01, 0.07)	0.13	-	-	-
Non-call related smartphone usage per day (hours)		-0.08	(-0.53, 0.37)	0.73	-	-	-
Ownership of smartphone (years)		0.34	(-0.36, 1.04)	0.33	-	-	-
Time from waking to first smartphone usage in the morning (minutes)		0.02	(-0.03, 0.080)	0.39	-	-	-
Number of additional devices owned		-0.45	(-3.29, 2.39)	0.75	-	-	-
Most personal smartphone feature							
Entertainment	19.92 (16.96)	4.16	(0.32, 8.00)	0.03*	4.21	(0.46, 7.96)	0.03*
Non entertainment	15.76 (13.83)						

^aSimple linear regression^bMultiple linear regression ($R^2 = 0.044$)^cCrude regression coefficient^dAdjusted regression coefficient

Notes: Multiple linear regression after controlling factors of sociodemography, socioeconomic status and academic achievement

*p value = statistically significant at <0.05

South Korean students, who used smartphones for 2–4 hours per day on average.¹⁵ In contrast, given the age disparities, a study conducted in Pakistan reported 11–14 hours daily smartphone used among medical practitioners (mean age, 23 years).¹⁶ Malaysians appear to be following local trends by spending more time on quick gratification applications like WhatsApp, Instagram, and Facebook.¹⁷ This era has a culture in which smartphones are their companions for playing, chatting, shopping, and spending their free time that is vastly different from that of their parents. The participants in the current study utilised their smartphones evenly for entertainment (50%) and non-entertainment (50%), which contrasts with a study from India, where smartphones were mostly used for social communication.¹⁸ We also discovered that participants used their smartphones 15 minutes after getting up in the morning, which differs from the majority of Swiss adolescents, who do so within 5 minutes.¹⁹ In light of the devices' affordability and availability, the number of devices possessed was slightly increased, i.e. having an extra device, comparable to that reported in a local research.¹⁷ According to the universal gratification theory (UGT), individuals resort to technology to fulfil their social and psychological needs makes it difficult to draw definite conclusions regarding smartphone use since cultures, values, and beliefs vary everywhere.²⁰

The majority of medical students in our study scored zero for self-reported symptoms, indicating that they did not have any (range 24.6–81.3%), which contradicts with our hypothesis that frequent smartphone usage may be associated with higher frequency of self-reported symptoms. This is further supported by an overall low TSS score of 17.84. In our study, 63.9% self-reported symptom of headache which is slightly more than that reported in Iranian students (53.5%) but less than that of Iraqi students, 82.1% of whom experienced headache.^{8,10}

This is nevertheless an issue, given the number of students who reported symptoms of varying severity, including insomnia (75.4%), neck discomfort (67.9%), headache (63.9%), visual difficulties (60.7%), and overall fatigue (56.7%). The distribution of symptoms differed from that of Iraqi students; in particular, insomnia (58.5%), neck pain (73.8%), headache (64.6%), visual difficulties (82.1%), and overall tiredness (69.9%).⁸ The National Radiological Protection Board addressed the concern in 2011 that a substantial proportion of frequency radiation may ultimately permeate the user's body parts and head, as well as up to two inches into the human brain.²¹ Current International Commission on Non-Ionizing Radiation Protection recommendations have highlighted concerns in adults, where smartphone radiation has a detrimental impact on cognitive functioning, cell communication, and fertility.²¹ In contrast, contrary findings show that smartphones pose no such hazards and are not strong enough to destroy atomic bonds or tissue.³ Because modern smartphones are more user-friendly, safer, and ergonomic than older smartphones or mobile phones, the present study's mean TSS score is low.³ The impact on human health is unknown since they are classed as category 2B by the International Agency for Research on Cancer (minimal evidence in humans but less than sufficient data in experimental animals).²² With years of cancer and neurodegenerative disease delays, it may take some time for the negative consequences to become apparent; by then, it may be too late.³

Further bivariate analysis indicated that being Malay, having working and highly educated parents, also academic achievement are important factors linked with smartphone usage. The Malay ethnicity was crucial since it was the predominant ethnic group in Malaysia. The findings on working parents contradict a Korean research, which found that the parents of the most smartphone addicted

children were mostly unemployed and had just a high school education.²³ The explanation for this might be that the sample group included children rather than university students. On the other hand, our results of increased smartphone usage outcomes contrast with those of Iranian students, who reported worse academic success because pupils were mentally occupied, resulting in a lack of concentration and focus in the lessons.²⁴ A local study also found highly educated parents had significant association with smartphone usage and certain smartphone patterns (making or receiving calls, purpose other than calls, extra devices owned), indicating high socioeconomic status of parents' ability to support students' monthly smartphone payments.²⁰ Aside from the desire for amusement, an American study found that when students utilise electronic educational devices, such as smartphones, throughout class sessions, their success improves by maximising time on task completion and using the camera to turn abstract concepts into tangible thoughts.²⁵ According to another research, smartphones increase students' productivity and educational achievement²⁵ by allowing them to download online lectures, read e-books, and use applications such as clouds to produce, post, retrieve, and exchange learning material with their friends.

The severity of symptoms reported by the participants in the current research was predicted by two major predictors. Firstly, the participants' TSS was influenced by their father's employment position. Our findings contradict a South Korean study, probably due to age differences, in which the study represented parents' employment insecurity and less than a high school degree.²³ Those parents are most likely ignorant of their responsibility to monitor their children's smartphone use and provide poor care for the child in terms of detecting the symptoms that their children may acquire as a result of smartphone exposure. It has also been established that trends

in children's smartphone use can result in psychological behavioural patterns²⁶: prospects for social interactions and physical activity may decrease as a result of smartphone fascination, resulting in negative physical symptoms in the long run. These findings indicate that smartphone usage has a detrimental impact on children's educational environments and development, which is evident in symptom severity.

We discovered that the smartphone's entertainment feature is another factor that contributes to higher TSS scores, which is consistent with an Iranian study.¹⁰ Participants in the current study indicated entertainment activities such as listening to music, playing games, watching movies, reading comic books, and maintaining social connection, particularly during the lockdown time. This is supported by Zuckerman's sensation-seeking theory of experience, which explains how people discover stimulus and resilience to dullness.²⁷ Prolonged smartphone use, on the other hand, may result in a variety of symptom severity due to: (a) poor ergonomics (headaches, neck, shoulder, and jaw pain) from anterior head syndrome¹ and (b) EMF or blue light exposure (insomnia, vision changes, difficulty concentrating, dry and irritated eyes).²⁸ The adoption of education and entertainment technologies, which cause excessive tension to the spine and digital eye strain, might have a severe impact on the younger generation who are still developing.⁸⁹

Our findings have certain limitations in terms of generalisation. To begin with, doing the study in different contexts was unavoidable. The research was first done prior to the countrywide COVID-19 lockdown order being issued⁶, and half of the results were received during the lockdown period, implying a lengthier screening time since each student relied significantly upon on e-learning system. We are unable to eliminate the possibility of reporting bias when using self-reported

data. Nonetheless, the study adds to our understanding of the impact of smartphone exposure on the severity of self-reported symptoms as perceived by participants, as well as previous evidence of smartphones influencing mental health.^{4, 5, 11, 26, 29}

CONCLUSION

During the early period of COVID-19 lockdown in Malaysia, frequency of medical students' self-reported symptoms were independently associated with their father's employment status and use of smartphone for entertainment purposes only. Despite this, the overall effect of smartphone usage on self-reported symptoms in general is low with the majority of medical students reporting no symptoms.

In the absence of regulation that controls healthy smartphone usage, parental involvement and supervision must begin in early education. Supportive systems that offer counselling or avoidance treatment on various forms of entertainment should be created. Public awareness initiatives for appropriate smartphone use beginning in early infancy are critical for reducing smartphone reliance. Parental smartphone usage and patterns may have a moderating influence. More research is required to establish a causal connection between smartphone usage and patterns and its association with other health-related outcomes, to understand the long-term implications. There is no clear evidence of harmful health impacts, yet no research has proved smartphones are safe, either.

ACKNOWLEDGEMENTS

We appreciate the participants' contributions and experiences. We would like to express our gratitude to the Dean of the Faculty of Medicine at UKM for granting us permission to publish this article.

FINANCIAL SUPPORT

The authors received no financial support for the research, authorship and/or publication of this article.

CONTRIBUTION

Both authors contributed equally, have read and approved the final version of the manuscript and agreed to be accountable for all aspects of the work.

CONFLICT OF INTEREST

The authors declared no potential conflicts of interest with respect to research, authorship, and/or publication of this article.

AVAILABILITY OF DATA AND MATERIALS

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

CONFERENCE PRESENTATION

This study has been presented at the 22nd National Public Health Colloquium, Department of Community Health, Universiti Kebangsaan Malaysia on May 4th-6th 2021, online event.

REFERENCES

- 1: Ali OTO. [C6-C7 and C7-T1 Facet Joint Osteoarthritis Cause By Prolonged Use of Smart Phone](#). Multi-Knowledge Electronic Comprehensive Journal For Education & Science Publications (MECSJ). 2019;26:1-9. [Accessed on 2022 April 10].
- 2: Cherry N. [Cell phone radiation poses a serious biological and health risk](#). Lincoln University, Canterbury, New Zealand, Environmental Management and Design Division;2001. [Accessed on 2022 April 10].
- 3: Mortazavi SMJ, Atefi M, Kholgh F. [The Pattern of Mobile Phone Use and Prevalence of Self-Reported Symptoms in Elementary and Junior High School Students in Shiraz, Iran](#). Iranian Journal of Medical Sciences. 2011;36:96-103. [Accessed on 2022 April 10].

- 4: Norbaidurah Ithnain, Shazli Ezzat Ghazali, Jaafar N. [Relationship between Smartphone Addiction with Anxiety and Depression among Undergraduate Students in Malaysia](#). International Journal of Health Sciences & Research. 2018;8:163-71. [Accessed on 2022 April 10].
- 5: Azlina Wati Nikmat, Nurul Azreen Hashim, Muhammad Farid Saidi, Nur Suhailah Mohd Zaki, Nur Nabihah Hasan Shukri, Abdulla NB. The Use And Addiction To Smart Phones Among Medical Students And Staffs In A Public University In Malaysia. ASEAN Journal of Psychiatry. 2018;19:1-7.
- 6: Eizairi Shamsudin. [14-day Movement Control Order begins nationwide on Wednesday](#). New Straits Times.[Internet] 2020 16 March 2020. [Accessed on 2022 April 10].
- 7: Kish L. [Survey Sampling](#). New York: John Wiley and Sons, Inc; 1965. [Accessed on 2022 April 10].
- 8: Sameeha Naser Abed, Rawaa Kamel Abd, Israa Dawood Salim, Nadia Abdul, Razzaq Jamal. [Health problems of Mobile Phone Addiction for sample of students and their health awareness at Institute Technical of Kut](#). Journal of Pharmaceutical Sciences and Research. 2018;10:412-5. [Accessed on 2022 April 10].
- 9: Ching SM, Yee A, Ramachandran V, Sazlyna Mohd Sazly Lim, Wan Aliaa Wan Sulaiman, Foo YL, et al. [Validation of a Malay Version of the Smartphone Addiction Scale among Medical Students in Malaysia](#). PLOS One. 2015;10:1-11. [Accessed on 2022 April 10].
- 10: Seyed Mohammad Javad Mortazavi, Mohammad Atefi, Fatemeh Kholghi. [The Pattern of Mobile Phone Use and Prevalence of Self-Reported Symptoms in Elementary and Junior High School Students in Shiraz, Iran](#). Iranian Journal of Medical Sciences. 2011;36:96-103. [Accessed on 2022 April 10].
- 11: Alhassan AA, Alqadhib EM, Taha NW, Alahmari RA, Salam M, Almutairi AF. [The relationship between addiction to smartphone usage and depression among adults: a cross sectional study](#). BMC Psychiatry. 2018; 18:148. [Accessed on 2022 April 10].
- 12: Demirci K, Akgonul M, Akpinar A. [Relationship of smartphone use severity with sleep quality, depression, and anxiety in university students](#). Journal of Behavioral Addictions. 2015;4:85-92. [Accessed on 2022 April 10].
- 13: Selvanathan M, Nur Atikah Mohamed Hussin, Noor Alyani Nor Azazi. [Students learning experiences during COVID-19: Work from home period in Malaysian Higher Learning Institutions](#). Teaching Public Administration. 2020:1-10. [Accessed on 2022 April 10].
- 14: Maurya P, Penuli Y, Kunwar A, Lalia H, Negi V, Williams A, et al. [Impact of Mobile Phone Usage on Psychosocial Wellbeing of Student Nurses](#). Journal of Nursing and Health Science. 2014; 3:39-42. [Accessed on 2022 April 10].
- 15: Lee KE, Si Heon Kim, Ha TY, Yoo YM, Han JJ, Jung JH, et al. [Dependency on Smartphone Use and Its Association with Anxiety in Korea](#). Public Health Reports. 2016;131:411-131. [Accessed on 2022 April 10].
- 16: Larik RSA, Mallah GA, Talpur MMA, Saeed A, Larik FA, Suhag AK, et al. [Adverse Effects of Cell Phone Radiation on Human Health](#). International Journal of Scientific & Engineering Research. 2016;7:480-6. [Accessed on 2022 April 10].
- 17: Kumar LR, Chii KD, Way LC, Jetly Y, Rajendaran V. [Awareness of mobile phone hazards among university students in a Malaysian medical school](#). Health. 2011;03:406-15. [Accessed on 2022 April 10].
- 18: Ganganahalli P, Tondare MB, Durgawale PM. [Use of Electronic Gadgets among Medical Students in Western Maharashtra, India](#). International Journal of Health Sciences and Research. 2014;4:26-30. [Accessed on 2022 April 10].
- 19: Haug S, Castro RP, Kwon M, Filler A, Kowatsch T, Schaub MP. [Smartphone use and smartphone addiction among young people in Switzerland](#). J Behav Addict. 2015;4:299-307. [Accessed on 2022 April 10].
- 20: Balakrishnan V, Raj RG. Exploring the relationship between urbanized Malaysian youth and their mobile phones: A quantitative approach. Telematics and Informatics. 2012;29:263-72.
- 21: International Commission on Non-Ionizing Radiation Protection. [Guidelines for Limiting Exposure to Time-Varying Electric and Magnetic Fields \(1 Hz to 100 kHz\)](#). Health Physics. 2010;99:818-36. [Accessed on 2022 April 10].
- 22: Kim KH, Kabir E, Jahan SA. The use of cell phone and insight into its potential human health impacts. Environmental, Monitoring and Assessment. 2016; 188:221.
- 23: Cho KS, Lee JM. [Influence of smartphone addiction proneness of young children on problematic behaviors and emotional intelligence: Mediating self-assessment effects of parents using smartphones](#). Computers in Human Be-

- havior. 2017;66:303-11. [Accessed on 2022 April 10].
- 24: Amidtaher M, Saadatmand MAS, Moghadam Z, Fathi G, Afshar R. [The Relationship between Mobile Cellphone Dependency, Mental Health and Academic Achievement](#). American Journal of Educational Research. 2016;4:408-11. [Accessed on 2022 April 10].
- 25: Woodcock B, Middleton A, Nortcliffe A. [Considering the Smartphone Learner: an investigation into student interest in the use of personal technology to enhance their learning](#). Student Engagement and Experience Journal. 2012;1:1-15. [Accessed on 2022 April 10].
- 26: Tan QYJ, Hartanto A, Toh WX, Yan H. [Commentary: Influence of smartphone addiction proneness of young children on problematic behaviors and emotional intelligence: Mediating self-assessment effects of parents using smartphones](#). Frontiers in Psychology. 2019;10:1-3. [Accessed on 2022 April 10].
- 27: Zuckerman M. Sensation seeking: beyond the optimal level of arousal. New York: Hillsdale, N.J., Erlbaum Associates; 1979.
- 28: Baeuchamp N. [4 Sneaky Ways Technology is affecting your family's health and may affect your child's new school year \[Internet\]](#). Exceptional Parent (EP) Magazine. 2019. [Accessed on 2022 April 10].
- 29: Xie H, Tao S, Zhang Y, Tao F, Wu X. [Impact of problematic mobile phone use and insufficient physical activity on depression symptoms: a college-based follow-up study](#). BMC Public Health. 2019; 19:1640. [Accessed on 2022 April 10].
-