



OFFICIAL PUBLICATION OF
THE MINISTRY OF HEALTH,
BRUNEI DARUSSALAM

Brunei International Medical Journal

Volume 15

30 April 2019 (24 Sya'ban 1440H)

BEHAVIOURAL HABITS AND ORAL CANCERS RISK AMONG THE YEMENI POPULATION: A CASE-CONTROL STUDY.

Azmawati Mohammed Naw¹, Hasanain Faisal Ghazi², Najla Mashlol³, Mohd Rohaizat Hassan¹,
Norfazilah Ahmad¹

¹Department of Community Health, Faculty of Medicine, Universiti Kebangsaan Malaysia.

²Community Medicine Unit, International Medical School, Management and Science University,
Selangor, Malaysia.

³ High Health Institution of Hodeidah, Yemen.

ABSTRACT

Aims: This study aimed to determine the association between behavioural habits and oral cancers risk among Yemeni population. **Methods:** A hospital-based unmatched case-control study was conducted among 74 oral cancers cases and 74 controls. Cases and controls were recruited using convenient sampling method. The data collection was performed using a guided questionnaire consists of socio-demographic characteristics and behavioural habits. Logistic regression analyses were conducted to determine the factors associated with oral cancers. **Results:** Older age and patients living in rural residence were significantly associated with oral cancers. Behavioural habits related to poor oral hygiene status [aOR=48.89 (95% CI: 5.45-408.26), $P < 0.001$] had the highest odds for oral cancers, followed by shammah consumption [aOR=3.34 (95% CI: 1.43-7.83), $P = 0.005$] among the Yemenis. **Conclusion:** Oral hygiene status and shammah consumption were modifiable behavioural habits that could be the focus for future health education and awareness programme strategies, especially among the older patients living in the rural residential area.

Keywords: Oral cancers, Oral hygiene, Tobacco usage, Habits, Case-Control

Brunei Int Med J. 2019;15:58-63

Brunei International Medical Journal (BIMJ)

Official Publication of the Ministry of Health, Brunei Darussalam

EDITORIAL BOARD

Editor-in-Chief	William Chee Fui CHONG
Sub-Editors	Vui Heng CHONG Ketan PANDE
Editorial Board Members	Nazar LUQMAN Muhd Syafiq ABDULLAH Alice Moi Ling YONG Ahmad Yazid ABDUL WAHAB Jackson Chee Seng TAN Dipo OLABUMUYI Pemasiri Upali TELISINGHE Roselina YAAKUB Pengiran Khairol Asmee PENGIRAN SABTU Dayangku Siti Nur Ashikin PENGIRAN TENGAH

INTERNATIONAL EDITORIAL BOARD MEMBERS

Lawrence HO Khok Yu (Singapore)	Surinderpal S BIRRING (United Kingdom)
Emily Felicia Jan Ee SHEN (Singapore)	Leslie GOH (United Kingdom)
John YAP (United Kingdom)	Chuen Neng LEE (Singapore)
Christopher HAYWARD (Australia)	Jimmy SO (Singapore)
Jose F LAPENA (Philippines)	Simon Peter FROSTICK (United Kingdom)

Advisor

Wilfred PEH (Singapore)

Past Editors

Nagamuttu RAVINDRANATHAN
Kenneth Yuh Yen KOK

Proof reader

John WOLSTENHOLME (CfBT Brunei Darussalam)

three relevant references should be included. Only images of high quality (at least 300dpi) will be acceptable.

Technical innovations

This section include papers looking at novel or new techniques that have been developed or introduced to the local setting. The text should not exceed 1000 words and should include not more than 10 figures illustration and references should not be more than 10.

Letters to the Editor

Letters discussing a recent article published in the BIMJ are welcome and should be sent to the Editorial Office by e-mail. The text should not exceed 250 words; have no more than one figure or table, and five references.

Criteria for manuscripts

Manuscripts submitted to the BIMJ should meet the following criteria: the content is original; the writing is clear; the study methods are appropriate; the data are valid; the conclusions are reasonable and supported by the data; the information is important; and the topic has general medical interest. Manuscripts will be accepted only if both their contents and style meet the standards required by the BIMJ.

Authorship information

Designate one corresponding author and provide a complete address, telephone and fax numbers, and e-mail address. The number of authors of each paper should not be more than twelve; a greater number requires justification. Authors may add a publishable footnote explaining order of authorship.

Group authorship

If authorship is attributed to a group (either solely or in addition to one or more individual authors), all members of the group must meet the full criteria and requirements for authorship described in the following paragraphs. One or more authors may take responsibility 'for' a group, in which case the other group members are not authors, but may be listed in an acknowledgement.

Authorship requirement

When the BIMJ accepts a paper for publication, authors will be asked to sign statements on (1) financial disclosure, (2) conflict of interest and (3) copyright transfer. The correspondence author may sign on behalf of co-authors.

Authorship criteria and responsibility

All authors must meet the following criteria: to have participated sufficiently in the work to take public responsibility for the content; to have made substantial contributions to the conception and de-

sign, and the analysis and interpretation of the data (where applicable); to have made substantial contributions to the writing or revision of the manuscript; and to have reviewed the final version of the submitted manuscript and approved it for publication. Authors will be asked to certify that their contribution represents valid work and that neither the manuscript nor one with substantially similar content under their authorship has been published or is being considered for publication elsewhere, except as described in an attachment. If requested, authors shall provide the data on which the manuscript is based for examination by the editors or their assignees.

Financial disclosure or conflict of interest

Any affiliation with or involvement in any organisation or entity with a direct financial interest in the subject matter or materials discussed in the manuscript should be disclosed in an attachment. Any financial or material support should be identified in the manuscript.

Copyright transfer

In consideration of the action of the BIMJ in reviewing and editing a submission, the author/s will transfer, assign, or otherwise convey all copyright ownership to the Clinical Research Unit, RIPAS Hospital, Ministry of Health in the event that such work is published by the BIMJ.

Acknowledgements

Only persons who have made substantial contributions but who do not fulfill the authorship criteria should be acknowledged.

Accepted manuscripts

Authors will be informed of acceptances and accepted manuscripts will be sent for copyediting. During copyediting, there may be some changes made to accommodate the style of journal format. Attempts will be made to ensure that the overall meaning of the texts are not altered. Authors will be informed by email of the estimated time of publication. Authors may be requested to provide raw data, especially those presented in graph such as bar charts or figures so that presentations can be constructed following the format and style of the journal. Proofs will be sent to authors to check for any mistakes made during copyediting. Authors are usually given 72 hours to return the proof. No response will be taken as no further corrections required. Corrections should be kept to a minimum. Otherwise, it may cause delay in publication.

Offprint

Contributors will not be given any offprint of their published articles. Contributors can obtain an electronic reprint from the journal website.

DISCLAIMER

All articles published, including editorials and letters, represent the opinion of the contributors and do not reflect the official view or policy of the Clinical Research Unit, the Ministry of Health or the institutions with which the contributors are affiliated to unless this is clearly stated. The appearance of advertisement does not necessarily constitute endorsement by the Clinical Research Unit or Ministry of Health, Brunei Darussalam. Furthermore, the publisher cannot accept responsibility for the correctness or accuracy of the advertisers' text and/or claim or any opinion expressed.

Aim and Scope of Brunei International Medical Journal

The Brunei International Medical Journal (BIMJ) is a six monthly peer reviewed official publication of the Ministry of Health under the auspices of the Clinical Research Unit, Ministry of Health, Brunei Darussalam.

The BIMJ publishes articles ranging from original research papers, review articles, medical practice papers, special reports, audits, case reports, images of interest, education and technical/innovation papers, editorials, commentaries and letters to the Editor. Topics of interest include all subjects that relate to clinical practice and research in all branches of medicine, basic and clinical including topics related to allied health care fields. The BIMJ welcomes manuscripts from contributors, but usually solicits reviews articles and special reports. Proposals for review papers can be sent to the Managing Editor directly. Please refer to the contact information of the Editorial Office.

Instruction to authors

Manuscript submissions

All manuscripts should be sent to the Managing Editor, BIMJ, Ministry of Health, Brunei Darussalam; e-mail: editor-in-chief@bimjonline.com. Subsequent correspondence between the BIMJ and authors will, as far as possible via should be conducted via email quoting the reference number.

Conditions

Submission of an article for consideration for publication implies the transfer of the copyright from the authors to the BIMJ upon acceptance. The final decision of acceptance rests with the Editor-in-Chief. All accepted papers become the permanent property of the BIMJ and may not be published elsewhere without written permission from the BIMJ.

Ethics

Ethical considerations will be taken into account in the assessment of papers that have experimental investigations of human or animal subjects. Authors should state clearly in the Materials and Methods section of the manuscript that institutional review board has approved the project. Those investigators without such review boards should ensure that the principles outlined in the Declaration of Helsinki have been followed.

Manuscript categories

Original articles

These include controlled trials, interventional studies, studies of screening and diagnostic tests, outcome studies, cost-effectiveness analyses, and large-scale epidemiological studies. Manuscript should include the following; introduction, materials and methods, results and conclusion. The objective should be stated clearly in the introduction. The text should not exceed 2500 words and references not more than 30.

Review articles

These are, in general, invited papers, but unsolicited reviews, if of good quality, may be considered. Reviews are systematic critical assessments of

literature and data sources pertaining to clinical topics, emphasising factors such as cause, diagnosis, prognosis, therapy, or prevention. Reviews should be made relevant to our local setting and preferably supported by local data. The text should not exceed 3000 words and references not more than 40.

Special Reports

This section usually consist of invited reports that have significant impact on healthcare practice and usually cover disease outbreaks, management guidelines or policy statement paper.

Audits

Audits of relevant topics generally follow the same format as original article and the text should not exceed 1,500 words and references not more than 20.

Case reports

Case reports should highlight interesting rare cases or provide good learning points. The text should not exceed 1000 words; the number of tables, figures, or both should not be more than two, and references should not be more than 15.

Education section

This section includes papers (i.e. how to interpret ECG or chest radiography) with particular aim of broadening knowledge or serve as revision materials. Papers will usually be invited but well written paper on relevant topics may be accepted. The text should not exceed 1500 words and should include not more than 15 figures illustration and references should not be more than 15.

Images of interest

These are papers presenting unique clinical encounters that are illustrated by photographs, radiographs, or other figures. Image of interest should include a brief description of the case and discussion with educational aspects. Alternatively, a mini quiz can be presented and answers will be posted in a different section of the publication. A maximum of

BEHAVIOURAL HABITS AND ORAL CANCERS RISK AMONG THE YEMENI POPULATION: A CASE-CONTROL STUDY.

Azmawati Mohammed Naw¹, Hasanain Faisal Ghazi², Najla Mashl³, Mohd Rohaizat Hassan¹, Norfazilah Ahmad¹

¹Department of Community Health, Faculty of Medicine, Universiti Kebangsaan Malaysia.

²Community Medicine Unit, International Medical School, Management and Science University, Selangor, Malaysia.

³ High Health Institution of Hodeidah, Yemen.

ABSTRACT

Aims: This study aimed to determine the association between behavioural habits and oral cancers risk among Yemeni population. **Methods:** A hospital-based unmatched case-control study was conducted among 74 oral cancers cases and 74 controls. Cases and controls were recruited using convenient sampling method. The data collection was performed using a guided questionnaire consists of socio-demographic characteristics and behavioural habits. Logistic regression analyses were conducted to determine the factors associated with oral cancers. **Results:** Older age and patients living in rural residence were significantly associated with oral cancers. Behavioural habits related to poor oral hygiene status [aOR=48.89 (95% CI: 5.45-408.26), $P<0.001$] had the highest odds for oral cancers, followed by shammah consumption [aOR=3.34 (95% CI: 1.43-7.83), $P = 0.005$] among the Yemenis. **Conclusion:** Oral hygiene status and shammah consumption were modifiable behavioural habits that could be the focus for future health education and awareness programme strategies, especially among the older patients living in the rural residential area.

Keywords: Oral cancers, Oral hygiene, Tobacco usage, Habits, Case-Control

INTRODUCTION

Globally, oral cancers has emerged as a significant cause of global public health concern.¹ The prevalence of oral cancers is particularly high among men and is the 8th most common cancer worldwide.¹ A total of 369,200 new cases of oral cancers were reported in 2012 with the majority of the cases were diagnosed in developing countries such as South and Southeast Asia.² The survival of oral cancers patients is poor (less than 50%

for 5-year survival rate) but with advancement in treatment such as chemotherapy and radiotherapy, the survival of oral cancers patients have been improved.^{3,4} The prevalence of oral cancers in Arab countries range from 1.8-2.13 per 100,000 populations with most of the patients diagnosed at the age of 50-60 years old.⁵ However, in Yemen, the prevalence was higher among the younger generation (less than 45 years old).⁵ According to the National Oncology Centre (NOC), oral cancers is the 5th most common cancer among the Yemenis.⁶ Behavioural habits such as smokeless tobacco (Shammah and khat) consumption and cigarette smoking were strongly associated with oral cancers.⁵ The hospital-based study also showed a very high preva-

Corresponding author: Azmawati Mohammed Naw¹ (M.Comm.Med), Dept. of Community Health, Faculty of Medicine, Universiti Kebangsaan Malaysia, Jln Yaacob Latiff, Bdr Tun Razak, Cheras, 56000 Kuala Lumpur, Malaysia.
Email: azmawati@ppukm.ukm.edu.my

prevalence of oral cancers in Yemen, where such behavioural habits are common.⁷ Hodeida city had the highest prevalence of oral cancers.⁶ However, there is limited information regarding the association of behavioural habits with oral cancers among the Yemenis in Hodeida city. Thus this study objective is to determine the associations between behavioural habits such as smokeless tobacco usage, smoking tobacco, habitual diet and oral hygiene status with oral cancers in Yemen.

MATERIALS AND METHODS

An unmatched case-control study was conducted among patients from two hospitals in the Hodeida governorate. The total sample size required for this study was 148 (74 cases and 74 controls) with 80% study power and at 5% significant level. The period for data collection was seven months from December 2013 to June 2014. The convenient sampling method was used to recruit the cases and controls. Cases were recruited from the Al-Amal Center of Cancer Treatment and controls were recruited from Althawra Hospital in Hodeida (approximately 2km apart). Cases with medically confirmed oral cancers were included, and cases with an unstable medical condition such as shortness of breath were excluded from the study. Controls with prior assessment and free from oral cancers or other oral-related cancers were included in this study. Controls with an unstable medical condition were also excluded from this study.

The data collection was conducted using a self-constructed and pre-tested questionnaire (See Appendix). After the informed consent taken from the respondents, a face to face guided interview done by trained interviewers, who were not blinded to the objectives of the study. Data were analysed by using the Statistical Package for Social Science (SPSS) version 20. Descriptive analyses were used to describe the categorical and

continuous variables, and Chi-square and t-test were performed to analyse the associations of patients' socio-demographic characteristics and oral cancers. Logistic regression analyses were conducted to obtain the crude odds ratio (cOR) and adjusted (aOR) and their 95% confidence interval (CI), as well as to determine the factors associated with oral cancers. The significance level in this study was set at $P < 0.05$.

RESULTS

Table 1 shows the mean age for the cases were significantly higher than the control group ($P = 0.003$). Percentages of female and not working patients among the cases was higher compared to control [(54.1% vs 50.0%) and (75.7% vs 63.5%), respectively]. There was a significantly higher percentage of patients living in the rural area among the cases (70.3%) compared to controls (41.9%), $P = 0.035$.

Table 2 depicts the behavioural habits of patients. For smokeless tobacco usage, the percentages of patients chewing khat and shammah usage were higher among the cases compared to controls. The mean total duration of shammah usage among cases was significantly longer than those for controls (25.35 vs 18.44, $p=0.01$). For tobacco smoking, cases who smoked cigarettes had a shorter total duration of smoking compared to control ($p<0.001$) but longer for cases who smoked shisha instead ($p=0.003$). Percentage distribution among the cases compared to control for alcohol, vegetables, fruits, spicy foods and hot drinks consumptions were almost balanced. However, there was a higher percentage of patients with poor oral hygiene status among the cases (compared to controls (98.6% vs 54.1% respectively, $p<0.001$).

Table 3 shows the factors associated with oral cancers. Based on multivariate regression analysis, only factors such as age

Table 1: Patients' socio-demographic characteristic and association with oral cancer.

Variables		Cases			Controls			p
		Mean (SD)	n	%	Mean (SD)	n	%	
Age in years		53.5 (16.4)			41.5 (18.2)			0.003 ^a
Gender	Male		34	45.9		37	50.0	NS
	female		40	54.1		37	50.0	
Occupation	Working		18	24.3		27	36.5	NS
	Not working		56	75.7		47	63.5	
Residential area	Rural		52	70.3		31	41.9	0.035 ^b
	Urban		22	29.7		43	58.1	
Education duration	<12years		7	9.5		19	25.7	NS
	≥12years		4	5.4		14	18.9	
	Not educated		63	85.1		41	55.4	
Family history of oral cancer	Yes		8	10.8		7	9.5	NS
	No		66	89.2		67	90.5	

^at-test; ^b chi-square test; NS = Non-significant, $p > 0.05$
Table 2: Patients' behavioural habits as modifiable risk factors for oral cancers, comparison between cases and controls.

Variables		Cases			Controls			p
		Mean(SD)	n	%	Mean(SD)	n	%	
Khat chewing	Yes		86.5	64		70.3	52	0.017
	No		13.5	10		29.7	22	
	Duration (years)	21.16(13.32)			17.21(11.66)			NS
	Frequency (week)	5.83(2.77)			4.98(2.65)			NS
	Duration of each session (hours)	3.73 (2.35)			3.23(2.09)			NS
Shammah usage	Yes		71.6	53		25.7	19	<0.001
	No		28.4	21		74.3	54	
	Duration (years)	25.35(14.90)			18.44(17.42)			0.010
	Frequency (day)	10.23(8.15)			6.72(3.97)			0.001
Cigarette smoking	Yes		16.2	12		17.6	13	NS
	No		83.8	62		82.4	61	
	Duration (years)	7.57(10.32)			16.38(12.21)			<0.001
	No of Cigarette	11.29(9.73)			13.92(11.69)			NS
Shisha smoking	Yes		47.3	35		37.8	28	NS
	No		52.7	39		62.2	46	
	Duration (years)	19.83(13.94)			13.59(10.86)			0.003
	Frequency (day)	2.20(1.84)			1.89(1.15)			NS
Alcohol drinking	Yes		1.4	1		0.0	0	NS
	No		98.6	73		100.0	74	
Vegetables consumption	Yes		100.0	74		98.6	73	NS
	No		0.0	0		1.4	1	
Fruits consumption	Yes		100.0	74		98.6	73	NS
	No		0.0	0		1.4	1	
Spicy food consumption	Yes		58.1	43		54.1	40	NS
	No		41.9	31		45.9	34	
Hot drinks consumption	Yes		48.6	36		47.3	35	NS
	No		51.4	38		52.7	39	
Oral hygiene status	Good		45.9	34		1.4	1	<0.001
	Poor		54.1	40		98.6	73	

Table 3: Modifiable and non-modifiable risk factors associated with oral cancer(s) among the Yemeni population.

Demographic Factors		cOR ^a	95% CI	p value	aOR ^b	95% CI	p value
Age (years)		1.04	1.02,1.06	<0.001	3.72	1.77,10.32	0.034
Gender	Female	1		NS	-	-	-
	Male	1.18	0.62,2.24		-	-	
Occupation	Not working	1		NS	-	-	-
	Working	1.79	0.88,3.64		-	-	
Education duration	<12 years	1		NS	-	-	-
	≥12 years	0.74	0.18,3.02		-	-	
	Not educated	3.86	1.48,10.04		-	-	
Residence	Urban	1		NS	1		0.004
	Rural	3.28	1.66,6.47		3.55	1.50,8.39	
Family history of oral cancer	No	1		NS	-	-	-
	Yes	1.16	0.40,3.38		-	-	
Behavioral habits		cOR ^a	95% CI	p value	aOR ^b	95% CI	p value
Khat chewing	No	1		0.026	-	-	-
	Yes	2.59	1.12,5.97		-	-	
Shammah usage	No	1		<0.001	1		0.005
	Yes	7.17	3.47,14.84		3.34	1.43,7.83	
Cigarette smoking	No	1		NS	-	-	-
	Yes	1.1	0.47,2.60		-	-	
Shisha Smoking	No	1		NS	-	-	-
	Yes	1.47	0.77,2.84		-	-	
Spicy food consumption	No	1		NS	-	-	-
	Yes	0.85	0.44,1.62		-	-	
Hot drinks consumption	No	1		NS	-	-	-
	Yes	0.95	0.50,1.81		-	-	
Oral hygiene status	Good	1		<0.001	1		<0.001
	Poor	0.02	0.01,0.12		48.89	5.85,408.26	

^a crude odds ratio (simple logistic regression); ^b adjusted odds ratio (multiple logistic regression)

[aOR=3.72 (95% CI 1.77,10.32), $p = 0.034$], residential area [aOR=3.55 (95% CI 1.50,8.39), $P = 0.004$], shammah usage [aOR=3.34 (95% CI 1.43,7.83), $p = 0.005$] and oral hygiene status [aOR=3.48.89 (95% CI 5.85,408.26), $p < 0.001$] were significant independent predictors for oral cancers.

DISCUSSION

This study showed that poor oral hygiene and smokeless tobacco consumption (shammah and khat usage), two modifiable behavioural habits, were independent predictors for development of oral cancers in Yemeni population in Hodeida city. Oral hygiene is essential for all individual to reduce incidence of oral disease. Poor oral hygiene may due to infre-

quent tooth brushing or using the chewing sticks.^{8, 9} Our findings on poor oral hygiene and the risk of oral cancers were in agreement with previous studies.^{8,10,11}

Our study confirmed that shammah usage increases the risk of developing oral cancers compared to control. Shammah is a preparation of a mixture of powdered tobacco, lime, ash, black pepper, oils and flavourings, which is frequently used in the Arabian Peninsula, especially in Saudi Arabia.¹² The use of shammah will increase the risk of oral leukoplakia-like lesions and could induce later development of oral cancers.^{13,14} Our study findings mirrored other similar studies carried out in middle-eastern Arab countries which showed a significant association of shammah

usage with increased risk of developing oral cancers.^{15,16}

Based on our findings, other factors not related to behavioural habits, such as age and geographical area of residence are significant risk factors for developing oral cancers. However, because we have recruited our controls from another hospital and did not match them for age or genders, these may have introduced sampling bias which may account for the significant findings. Despite this, several studies have previously reported that increasing age is associated with higher risk of developing oral cancers reflecting our study findings which showed the risk of developing oral cancers increases by 3.72 times for every year increase in age.^{17,18} Similarly since controls were recruited from Althawra Hospital, which is located in the city center while the Al-Amal center for cancer treatment is situated in the outskirts of the city, this may have influence the catchment of mainly rural patients as cases compare to the controls. Furthermore, patients from the Althawra hospital, coming from the city center may have better oral hygiene education compare to the cases from the rural area. The knowledge of oral cancers patients in the rural area especially regarding risk behaviour and screening methods has been shown to be low.¹⁹

There were few limitations in our study. The unmatched case-control design exposed the study to sampling bias as mentioned above. Data collections through interview lead to recall bias. In this case, the cases tend to report more on usage or exposure. In the process interview, we did not blind the interviewers of the study objective. This may lead to possibility of interviewer bias in this study. Despite the study limitations, the results could serve as baseline data for future prospective studies.

CONCLUSION

Both non-modifiable (Age and rural residence) and modifiable factors (behavioural habits such as poor oral hygiene and smokeless tobacco consumption) have a significant association with oral cancers. The modifiable behavioural habits above could form focus for future health education and awareness programme strategies, especially among the older patients living in the rural residential area.

ACKNOWLEDGEMENT

The authors would like to acknowledge our deepest appreciation and gratitude to the invaluable help in the study processes to the data collectors, all workers in the National Oncology Centre in Hodeida and all the patients who participated in this study.

AUTHOR'S CONTRIBUTIONS

All authors contributed to the processes of writing, revising editing of the paper and approved the final version for publication.

FUNDING

This study did not receive direct funding.

COMPETING INTEREST

All authors declare no conflicts of interest.

REFERENCES

- 1: Petersen PE. Strengthening the prevention of oral cancers: the WHO perspective. *Community Dent Oral Epidemiol* 2005;33: 397-399.
- 2: Ghantous Y1, Abu Elnaaj. Global incidence and risk factors of oral cancers. *Harefuah*. 2017;156(10):645-649.
- 3: Baykul T, Yilmaz HH, Aydin U, et al. Early diagnosis of oral cancers. *J Int Med Res*. 2010; 38:737-749. [Accessed on 2019 April 26]. Pdf available at <https://journals.sagepub.com/doi/pdf/10.1177/147323001003800302>
- 4: Cheraghloo S, Schettino A, Zogg CK, et al. Changing prognosis of oral cancers: An analysis of survival and treatment between 1973 and 2014. *Laryngoscope*. 2018;128(12):2762-2769.
- 5: Al-Jaber Abeer, Lubna Al-Nasser, Ashraf El-Metwally. Epidemiology of oral cancers in Arab countries. *Saudi Medical Journal*, 2016;37(3):249-255. [Accessed on 2019 April 26]. Pdf available at <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4800887/pdf/SaudiMedJ-37-249.pdf>

- 6: AL-Nabhi A, Ahmed MTA, Abdul Hamid G. Pattern of cancer in Yemen: first result from the national oncology center, Sana'a, 2007. *European Journal of Pharmaceutical and Medical Research.* 2017;4(1):149-154. [Accessed on 2019 April 26]. Pdf available at <https://pdfs.semanticscholar.org/b503/18c852dd9aa07e060ec5ab3264685b2d558c.pdf>
 - 7: Sawair FA, Al-Mutwakel A, Al-Eryani K, et al. High relative frequency of oral squamous cell carcinoma in Yemen: Khat and tobacco chewing as its aetiological background. *Int J Environ Health Res.* 2007;17:185-95.
 - 8: Oji C, Chukwuneke F. Poor oral Hygiene may be the Sole Cause of Oral cancers. *J Maxillofac Oral Surg.* 2012;11(4):379-383. [Accessed on 2019 April 26]. Pdf available at https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3485455/pdf/12663_2012_Article_359.pdf
 - 9: Macigo FG, Gathece LW, Guthua SW, et al. Oral hygiene practices and risk of oral leukoplakia. *East African Medical Journal* 2006;83(4):73-78. [Accessed on 2019 April 26]. Pdf available at <https://pdfs.semanticscholar.org/3204/ecb3251f-be4117a64e29e4449a37dd6d5733.pdf>
 - 10: Zheng TZ, Boyle P, Hu HF, et al. Dentition, oral hygiene, and risk of oral cancers: a case-control study in Beijing, People's Republic of China. *Cancer Causes Control.* 1990;1(3):235-241.
 - 11: Gupta B, Bray F, Kumar N, et al. Associations between oral hygiene habits, diet, tobacco and alcohol and risk of oral cancers: A case-control study from India. *Cancer Epidemiol.* 2017; 51:7-14.
 - 12: Bakdash A. Shammah (Smokeless Tobacco) and Public Health. *Asian Pac J Cancer Prev.* 2017;18(5):1183-1190. [Accessed on 2019 April 26]. Pdf available at <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC555521/pdf/APJCP-18-1183.pdf>
 - 13: Al-Tayar BA Tin-Oo MM, Sinor MZ, et al. Association between Shammah Use and Oral Leukoplakia-like Lesions among Adult Males in Dawan Valley, Yemen. *Asian Pac J Cancer Prev.* 2015;16(18):8365-8370.
 - 14: Kil TJ, Kim HS, Kim HJ, et al. Genetic Abnormalities in Oral Leukoplakia and Oral cancers Progression. *Asian Pac J Cancer Prev.* 2016;17(6):3001-3006. [Accessed on 2019 April 26]. Pdf available at http://journal.waocp.org/article_16383_5abcbc5500f79d26c8edc9f27895ef9e.pdf
 - 15: Idris AM, Vani NV, Saleh S, et al. Relative Frequency of Oral Malignancies and Oral Precancer in the Biopsy Service of Jazan Province, 2009-2014. *Asian Pac J Cancer Prev.* 2016;17(2):519-525. [Accessed on 2019 April 26]. Pdf available at http://journal.waocp.org/article_31879_698b6c083fe2e8cf9ed3490206d5a3a7.pdf
 - 16: Al-zamzami AA, Mohammed Suleiman A. Oral cancers among yemenis patient: A prospective hospital-based study. *Dent Med Res* 2018;6:32-36. [Accessed on 2019 April 26]. Pdf available at http://www.dmrjournal.org/temp/DentMedRes6232-3279066_090630.pdf
 - 17: Hari R, Jayanta S, Hemant K, et al. Oral cancers: Risk Factors and Molecular Pathogenesis. *J Maxillofac Oral Surg.* 2011;10(2):132-137. [Accessed on 2019 April 26]. Pdf available at https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3177522/pdf/12663_2011_Article_195.pdf
 - 18: Troeltzsch M, Knosel T, Eichinger C, et al. Clinicopathologic Features of Oral Squamous Cell Carcinoma: Do They Vary in Different Age Groups? *J Oral Maxillofac Surg* 2014;72:1291-1300. [Accessed on 2019 April 26]. Pdf available at [https://www.joms.org/article/S0278-2391\(14\)00101-3/pdf](https://www.joms.org/article/S0278-2391(14)00101-3/pdf)
 - 19: Riley JL, Pomery EA, Dodd VJ, et al. Disparities in Knowledge of Mouth or Throat Cancer Among Rural Floridians. *J Rural Health.* 2013;29(3):10.1111/jrh.12003. [Accessed on 2019 April 26]. Pdf available at <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3695415/pdf/nihms-455696.pdf>
-