

OFFICIAL PUBLICATION OF
THE MINISTRY OF HEALTH,
BRUNEI DARUSSALAM

Brunei International Medical Journal

Volume 16

13 May 2020 (20 Ramadhan 1441H)

PREVALENCE OF CRYPTO-SPORIDIOSIS AND ASSOCIATED RISK FACTORS AMONG PATIENTS WITH ACUTE DIARRHOEA ATTENDING ABUJA TEACHING HOSPITAL, NIGERIA.

Idris Nasir ABDULLAHI, Anthony Uchenna EMERIBE, Kabir UMAR, Hafeez Aderinsayo ADEKOLA, Peter Elisha GHAMBA, Nafiu FARUKU, Nkechi Blessing ONUKEGBE, Chisom Emmanuel OKECHUKWU, Adamu BABAYO, Dele Ohinoyi AMADU.

ABSTRACT

Background: Cryptosporidiosis is a zoonotic and foodborne parasitic infection which has become a major subject of public health concern in developing countries. This study aimed to determine the prevalence and associated risk factors of cryptosporidiosis in patients with acute diarrhea (<5 days post onset) attending University of Abuja Teaching Hospital (UATH), Abuja, Nigeria. **Materials and Methods:** 160 patients presenting with acute diarrhea (40 <5 years children, 40 HIV patients, 40 patients with active pulmonary tuberculosis (pTB) and 40 persons with type-2 diabetic mellitus (T2DM)). Fresh fecal samples were collected and immediately analyzed for *Cryptosporidium* spp using modified Ziehl Neelsen (mZN) protocol and *Cryptosporidium* antigen (CA) based-ELISA. **Results:** The detection rate of CA by ELISA was 26.3% while *Cryptosporidium* spp oocytes detection by mZN was 13.7%. The prevalence of cryptosporidiosis was 7.5% in pTB infected patients, 20% in HIV patients, 62.5% in <5 years children and 27.5% in T2DM patients. The prevalence of cryptosporidiosis was significantly associated with gender, socioeconomic status and marital status ($p < 0.05$) Possession of house pets (OR=9.21, 95% CI=4.15-20.4), practice of backyard poultry (OR=8.81, 95% CI=3.27-23.73), daily vegetable consumption (OR=4.42, 95% CI=2.02-9.63) and household size (OR= 2.49, 95% CI=1.14-5.43) were significant risk factors of cryptosporidiosis. **Conclusion:** Compared to the 15% national prevalence of cryptosporidiosis in Nigeria, it can be inferred that the present study revealed a significantly high prevalence of cryptosporidiosis among paediatric patients, HIV patients and T2DM patients presenting with acute diarrhea to Abuja Teaching Hospital.

Keywords: Cryptosporidiosis, Children, Type 2 Diabetes Mellitus, Diarrhea, HIV, Intestinal parasite, Risk factors, Tuberculosis.

Brunei Int Med J. 2020;16:49-57

Brunei International Medical Journal (BIMJ)

Official Publication of the Ministry of Health, Brunei Darussalam

EDITORIAL BOARD

Editor-in-Chief	William Chee Fui CHONG
Sub-Editors	Vui Heng CHONG Ketan PANDE
Editorial Board Members	Muhd Syafiq ABDULLAH Alice Moi Ling YONG Ahmad Yazid ABDUL WAHAB Jackson Chee Seng TAN Pemasiri Upali TELISINGHE Roselina YAAKUB Pengiran Khairol Asmee PENGIRAN SABTU Dayangku Siti Nur Ashikin PENGIRAN TENGAH

INTERNATIONAL EDITORIAL BOARD MEMBERS

Lawrence HO Khek Yu (Singapore)	Surinderpal S BIRRING (United Kingdom)
Emily Felicia Jan Ee SHEN (Singapore)	Leslie GOH (United Kingdom)
John YAP (United Kingdom)	Chuen Neng LEE (Singapore)
Christopher HAYWARD (Australia)	Jimmy SO (Singapore)
Jose F LAPENA (Philippines)	Nazar LUQMAN
Dipo OLABUMUYI	

Advisor

Wilfred PEH (Singapore)

Past Editors

Nagamuttu RAVINDRANATHAN
Kenneth Yuh Yen KOK

Proof reader

John WOLSTENHOLME (CfBT Brunei Darussalam)

three relevant references should be included. Only images of high quality (at least 300dpi) will be acceptable.

Technical innovations

This section include papers looking at novel or new techniques that have been developed or introduced to the local setting. The text should not exceed 1000 words and should include not more than 10 figures illustration and references should not be more than 10.

Letters to the Editor

Letters discussing a recent article published in the BIMJ are welcome and should be sent to the Editorial Office by e-mail. The text should not exceed 250 words; have no more than one figure or table, and five references.

Criteria for manuscripts

Manuscripts submitted to the BIMJ should meet the following criteria: the content is original; the writing is clear; the study methods are appropriate; the data are valid; the conclusions are reasonable and supported by the data; the information is important; and the topic has general medical interest. Manuscripts will be accepted only if both their contents and style meet the standards required by the BIMJ.

Authorship information

Designate one corresponding author and provide a complete address, telephone and fax numbers, and e-mail address. The number of authors of each paper should not be more than twelve; a greater number requires justification. Authors may add a publishable footnote explaining order of authorship.

Group authorship

If authorship is attributed to a group (either solely or in addition to one or more individual authors), all members of the group must meet the full criteria and requirements for authorship described in the following paragraphs. One or more authors may take responsibility 'for' a group, in which case the other group members are not authors, but may be listed in an acknowledgement.

Authorship requirement

When the BIMJ accepts a paper for publication, authors will be asked to sign statements on (1) financial disclosure, (2) conflict of interest and (3) copyright transfer. The correspondence author may sign on behalf of co-authors.

Authorship criteria and responsibility

All authors must meet the following criteria: to have participated sufficiently in the work to take public responsibility for the content; to have made substantial contributions to the conception and de-

sign, and the analysis and interpretation of the data (where applicable); to have made substantial contributions to the writing or revision of the manuscript; and to have reviewed the final version of the submitted manuscript and approved it for publication. Authors will be asked to certify that their contribution represents valid work and that neither the manuscript nor one with substantially similar content under their authorship has been published or is being considered for publication elsewhere, except as described in an attachment. If requested, authors shall provide the data on which the manuscript is based for examination by the editors or their assignees.

Financial disclosure or conflict of interest

Any affiliation with or involvement in any organisation or entity with a direct financial interest in the subject matter or materials discussed in the manuscript should be disclosed in an attachment. Any financial or material support should be identified in the manuscript.

Copyright transfer

In consideration of the action of the BIMJ in reviewing and editing a submission, the author/s will transfer, assign, or otherwise convey all copyright ownership to the Clinical Research Unit, RIPAS Hospital, Ministry of Health in the event that such work is published by the BIMJ.

Acknowledgements

Only persons who have made substantial contributions but who do not fulfill the authorship criteria should be acknowledged.

Accepted manuscripts

Authors will be informed of acceptances and accepted manuscripts will be sent for copyediting. During copyediting, there may be some changes made to accommodate the style of journal format. Attempts will be made to ensure that the overall meaning of the texts are not altered. Authors will be informed by email of the estimated time of publication. Authors may be requested to provide raw data, especially those presented in graph such as bar charts or figures so that presentations can be constructed following the format and style of the journal. Proofs will be sent to authors to check for any mistakes made during copyediting. Authors are usually given 72 hours to return the proof. No response will be taken as no further corrections required. Corrections should be kept to a minimum. Otherwise, it may cause delay in publication.

Offprint

Contributors will not be given any offprint of their published articles. Contributors can obtain an electronic reprint from the journal website.

DISCLAIMER

All articles published, including editorials and letters, represent the opinion of the contributors and do not reflect the official view or policy of the Clinical Research Unit, the Ministry of Health or the institutions with which the contributors are affiliated to unless this is clearly stated. The appearance of advertisement does not necessarily constitute endorsement by the Clinical Research Unit or Ministry of Health, Brunei Darussalam. Furthermore, the publisher cannot accept responsibility for the correctness or accuracy of the advertisers' text and/or claim or any opinion expressed.

Aim and Scope of Brunei International Medical Journal

The Brunei International Medical Journal (BIMJ) is a six monthly peer reviewed official publication of the Ministry of Health under the auspices of the Clinical Research Unit, Ministry of Health, Brunei Darussalam.

The BIMJ publishes articles ranging from original research papers, review articles, medical practice papers, special reports, audits, case reports, images of interest, education and technical/innovation papers, editorials, commentaries and letters to the Editor. Topics of interest include all subjects that relate to clinical practice and research in all branches of medicine, basic and clinical including topics related to allied health care fields. The BIMJ welcomes manuscripts from contributors, but usually solicits reviews articles and special reports. Proposals for review papers can be sent to the Managing Editor directly. Please refer to the contact information of the Editorial Office.

Instruction to authors

Manuscript submissions

All manuscripts should be sent to the Managing Editor, BIMJ, Ministry of Health, Brunei Darussalam; e-mail: editor-in-chief@bimjonline.com. Subsequent correspondence between the BIMJ and authors will, as far as possible via should be conducted via email quoting the reference number.

Conditions

Submission of an article for consideration for publication implies the transfer of the copyright from the authors to the BIMJ upon acceptance. The final decision of acceptance rests with the Editor-in-Chief. All accepted papers become the permanent property of the BIMJ and may not be published elsewhere without written permission from the BIMJ.

Ethics

Ethical considerations will be taken into account in the assessment of papers that have experimental investigations of human or animal subjects. Authors should state clearly in the Materials and Methods section of the manuscript that institutional review board has approved the project. Those investigators without such review boards should ensure that the principles outlined in the Declaration of Helsinki have been followed.

Manuscript categories

Original articles

These include controlled trials, interventional studies, studies of screening and diagnostic tests, outcome studies, cost-effectiveness analyses, and large-scale epidemiological studies. Manuscript should include the following; introduction, materials and methods, results and conclusion. The objective should be stated clearly in the introduction. The text should not exceed 2500 words and references not more than 30.

Review articles

These are, in general, invited papers, but unsolicited reviews, if of good quality, may be considered. Reviews are systematic critical assessments of

literature and data sources pertaining to clinical topics, emphasising factors such as cause, diagnosis, prognosis, therapy, or prevention. Reviews should be made relevant to our local setting and preferably supported by local data. The text should not exceed 3000 words and references not more than 40.

Special Reports

This section usually consist of invited reports that have significant impact on healthcare practice and usually cover disease outbreaks, management guidelines or policy statement paper.

Audits

Audits of relevant topics generally follow the same format as original article and the text should not exceed 1,500 words and references not more than 20.

Case reports

Case reports should highlight interesting rare cases or provide good learning points. The text should not exceed 1000 words; the number of tables, figures, or both should not be more than two, and references should not be more than 15.

Education section

This section includes papers (i.e. how to interpret ECG or chest radiography) with particular aim of broadening knowledge or serve as revision materials. Papers will usually be invited but well written paper on relevant topics may be accepted. The text should not exceed 1500 words and should include not more than 15 figures illustration and references should not be more than 15.

Images of interest

These are papers presenting unique clinical encounters that are illustrated by photographs, radiographs, or other figures. Image of interest should include a brief description of the case and discussion with educational aspects. Alternatively, a mini quiz can be presented and answers will be posted in a different section of the publication. A maximum of

PREVALENCE OF CRYPTO-SPORIDIOSIS AND ASSOCIATED RISK FACTORS AMONG PATIENTS WITH ACUTE DIARRHOEA ATTENDING ABUJA TEACHING HOSPITAL, NIGERIA.

Idris Nasir ABDULLAHI¹, Anthony Uchenna EMERIBE², Kabir UMAR¹, Hafeez Aderinsayo ADEKOLA³, Peter Elisha GHAMBA⁴, Nafiu FARUKU¹, Nkechi Blessing ONUKEGBE⁵, Chisom Emmanuel OKECHUKWU⁶, Adamu BABAYO⁷, Dele Ohinoyi AMADU⁸

¹Department of Medical Laboratory Science, Faculty of Allied Health Sciences, Ahmadu Bello University, Zaria, Nigeria, ²Department of Medical Laboratory Science, Faculty of Allied Health Sciences, University of Calabar, Calabar, Nigeria, ³Department of Microbiology, Olabisi Onabanjo University, Ogun State, Nigeria, ⁴WHO National Polio Laboratory, University of Maiduguri Teaching Hospital, Maiduguri, Nigeria, ⁵Department of Strategic Information and Research, Institute of Human Virology, Abuja, Nigeria, ⁶Department of Medical Microbiology, School of Medical Laboratory Science, Usmanu Danfodiyo University, Sokoto, Nigeria, ⁷Department of Medical Microbiology and Parasitology, Faculty of Clinical Sciences, Bayero University, Kano, Nigeria, and ⁸Department of Medical Microbiology and Parasitology, University of Ilorin Teaching Hospital, Ilorin, Nigeria.

ABSTRACT

Background: Cryptosporidiosis is a zoonotic and foodborne parasitic infection which has become a major subject of public health concern in developing countries. This study aimed to determine the prevalence and associated risk factors of cryptosporidiosis in patients with acute diarrhea (<5 days post onset) attending University of Abuja Teaching Hospital (UATH), Abuja, Nigeria. **Materials and Methods:** 160 patients presenting with acute diarrhea (40 <5 years children, 40 HIV patients, 40 patients with active pulmonary tuberculosis (pTB) and 40 persons with type-2 diabetic mellitus (T2DM)). Fresh fecal samples were collected and immediately analyzed for *Cryptosporidium spp* using modified Ziehl Neelsen (mZN) protocol and Cryptosporidium antigen (CA) based-ELISA. **Results:** The detection rate of CA by ELISA was 26.3% while *Cryptosporidium spp* oocytes detection by mZN was 13.7%. The prevalence of cryptosporidiosis was 7.5% in pTB infected patients, 20% in HIV patients, 62.5% in <5 years children and 27.5% in T2DM patients. The prevalence of cryptosporidiosis was significantly associated with gender, socioeconomic status and marital status ($p < 0.05$) Possession of house pets (OR=9.21, 95% CI=4.15-20.4), practice of backyard poultry (OR=8.81, 95% CI=3.27-23.73), daily vegetable consumption (OR=4.42, 95% CI=2.02-9.63) and household size (OR= 2.49, 95% CI=1.14-5.43) were significant risk factors of cryptosporidiosis. **Conclusion:** Compared to the 15% national prevalence of cryptosporidiosis in Nigeria, it can be inferred that the present study revealed a significantly high prevalence of cryptosporidiosis among paediatric patients, HIV patients and T2DM patients presenting with acute diarrhea to Abuja Teaching Hospital.

Keywords: Cryptosporidiosis, Children, Type 2 Diabetes Mellitus, Diarrhea, HIV, Intestinal parasite, Risk factors, Tuberculosis.

Corresponding author: Mr. Idris Nasir ABDULLAHI, Department of Medical Laboratory Science, Faculty of Allied Health Sciences, Ahmadu Bello University, Zaria, Nigeria
Email: eedris888@yahoo.com
Tel.: +2348027771068

INTRODUCTION

Cryptosporidium is an apicomplexan parasite, which is common among domestic and wild animals.¹ Transmission to humans occur mainly through fecal-oral route when they consume food and water contaminated with oocysts, or during recreational events (such as swimming pools).^{1,2} On several occasions cryptosporidiosis has been implicated in diarrhea outbreaks in many countries.²

Cryptosporidiosis often manifests as self-limiting diarrhea in immunocompetent persons, and as a progressively life-threatening diarrhea in people who are immunocompromised. These include persons with human immunodeficiency virus infection (HIV), active pulmonary tuberculosis (pTB), younger children and the elderly, persons on cancer chemotherapy, and any other condition that could compromise the immune function such as malnutrition.³

There are several studies which reported disproportionate prevalence of cryptosporidiosis in Nigeria. However, a recent meta-analysis revealed a national prevalence of cryptosporidiosis of 15% in Nigeria.⁴ Of these studies, few used enzyme immunoassays and compared results with that of the modified Zeihl Neelsen (mZN) technique. In addition, none simultaneously compared the prevalence of cryptosporidiosis across different study population. Hence, this study aims to determine the prevalence and associated risk factors of cryptosporidiosis among various categories of immunocompromised patients presenting with acute diarrhea at University of Abuja Teaching Hospital, Gwagwalada, Abuja, Nigeria.

MATERIALS AND METHODS

This hospital-based cross-sectional study was conducted at the University of Abuja Teaching Hospital, Gwagwalada, located at a suburb of Abuja, the federal capital territory (FCT),

Abuja, the federal capital territory (FCT), Abuja, Nigeria. According to the 2016 Nigerian National Population census, less than 10% of Gwagwalada area council had access to portable tap water. Ethical approval was obtained from the Human Ethical Research Committee (HREC) of UATH. The study was explained to patients or parents/guardians of children, and were enrolled into the study upon giving written and / or verbal informed consent. Structured questionnaire was used to collate socio-demographic variables and possible cryptosporidiosis risk factors from patients or parents/guardians. This was conducted in accordance with the Declaration of Helsinki. Data generated were anonymously analyzed throughout the study.

Between 2nd April to 30th November, 2018, patients were randomly recruited based on the following inclusion criteria; patients who have active pTB, HIV, T2DM or less than 5 years, presenting with acute diarrhea, and tested negative for intestinal parasites by routine wet preparation microscopy and have not started anti-protozoan therapy. However, those that who did not provide consent to participate in the study were excluded.

Interviewer-administered questionnaire was used to assess sociodemographic variables of participants, such as age, occupation, education level, civil status, monthly income, household, conditions, water availability, supply, washing fruit and vegetables. Indices of socioeconomic status (SES) was defined by educational attainment, income, housing, and employment variables into 3 groups: Upper SES, Middle SES and Lower SES (See APPENDIX I for full definition of indices of socioeconomic status).

Fresh diarrheic stool samples uncontaminated with urine were collected in a properly labeled (with patient's study number), leak proof, clean, sterile plastic containers with screw cap lid from patients. Samples

were transported to the laboratory and safety precautions were observed throughout the period of processing the specimens. Fresh samples were kept at 2 - 8° C and tested within 24 hours of collection.

A direct sample of the stool was made with both saline and iodine mounts on clean grease free slides and examined under the microscope first with 10× and then 40× objective lens for ova and cysts of parasites. A sample of the stool was concentrated using formol ether method concentration technique.⁵ Detection of *Cryptosporidium* oocysts in the concentrated stool was done using the modified cold Ziehl Neelsen staining technique, as previously described (See Appendix II).⁵ Determination of Fecal Cryptosporidia Antigen was also performed by Sandwich ELISA using Para-Tech® ELISA kit from *Medical Chemical Corp, California, USA* (See APPENDIX III).

The study sample size was determined using fishers' expression. Using the 4.8% prevalence of cryptosporidiosis from a previous cross-sectional study in North-central Nigeria, a minimum sample size of 72 was calculated.⁶ However, the number was increased to 160 (comprising of 40 <5years children, 40 HIV patients, 40 patients with active pTB and 40 patients with T2DM) to improve the statistical credence of the study. Data were presented as mean +/- SD and percentages. Categorical variables were compared with prevalence of cryptosporidiosis using chi-square test. Data was submitted for univariate logistic regression analyses to compute Odd Ratios (OR) at 95% confidence interval (95%CI) as a measure of risk factor for cryptosporidiosis. The various tests were carried out as two-tailed and outcomes with probability value below 0.05 were considered to be statistically significant.

RESULTS

The detection rate of Cryptosporidiosis by ELISA was 42/160 (26.3%) while *Cryptosporidium spp* was detected by mZN in 22/160 (13.7%) (Figure 1). The prevalence of cryptosporidiosis was 7.5% (3/40) in pTB patients, 20% (8/40) in HIV infected patients, 62.5% (25/40) in <5 years children and 27.5% in T2DM patients (Table I). There was significant association between prevalence of cryptosporidiosis with the various patients' groups ($\chi^2=32.15$; $p<0.0001$).

Children ≤5 years had the highest prevalence of cryptosporidiosis, 25 (62.5%) followed by those between 35 – 44 years, 3 (21.4%). However, none of those between 45 – 54 years had cases of cryptosporidiosis. The prevalence of cryptosporidiosis was relatively higher in males, 27(37.5%) than the female counterpart, 15 (17.0%). Patients without formal education had the highest prevalence of cryptosporidiosis, 22 (33.8%) and least among those with tertiary education, 3 (12.5%). The prevalence of cryptosporidiosis was highest among patients who were married, 27 (37.5%), followed by the singles, 14 (18.4%) and least in widows/ widowers, 1 (8.3%).

In regards to the residential areas of patients, rural residents had the highest prevalence of cryptosporidiosis, 21 (45.6%), followed by those who reside in urban communities, 8 (24.2%), and least in suburb settlers, 13 (16.0%). Patients who were unemployed had the highest prevalence of cryptosporidiosis, 12 (38.7%) and least among civil servants, 3 (10.3%). Patients who were at lower socioeconomic category had the highest prevalence of cryptosporidiosis, 29 (39.7%), and least in those with high socioeconomic status, 3 (7.3%). The prevalence of cryptosporidiosis was significantly associated with sex, socioeconomic status and marital status ($p< 0.05$), but not with education level, occupation, age and residential area ($p>0.05$) (Table II).




Figure 1: Cryptosporidium detection rates of modified Zeihl Neelsen stain and ELISA.

In regards to the risk factors of cryptosporidiosis, patients who had pets in their houses had relatively higher cases of cryptosporidiosis, 29 (55.8%) than those who did not, 13 (12.0%). Those that consumed vegetable foods on daily basis had relatively higher cases of cryptosporidiosis, 31 (40.2%) than who did not, 11 (13.3%). Patients who had boreholes/ wells as their sole sources of drinking water had relatively higher cases of cryptosporidiosis, 23 (27.1) than those who outsourced their drinking water from taps, 19 (25.3%). Persons who practiced solely open bush defecation had higher cases of cryptosporidiosis, 22 (32.4%) than those who defecate using only pit latrines and water closet, 20 (21.7%). Patients who resided in a densely populated households had higher cases of

cryptosporidiosis, 31 (33.7%) than those who resided in sparsely populated houses, 11 (16.9%). Persons who had backyard poultry in their houses had higher cases of cryptosporidiosis, 15 (68.2%) than those who do not, 27 (19.6%).

After univariate logistic regression, possession of house pets (OR=9.21, 95% CI=4.15-20.4), practice of backyard poultry (OR=8.81, 95% CI=3.27-23.73), daily vegetable consumption (OR=4.42, 95% CI=2.02-9.63) and household size (OR= 2.49, 95% CI=1.14-5.43) were significant risk factors of cryptosporidiosis (Table III).

DISCUSSION

Cryptosporidiosis, which is rarely seen in immunocompetent individuals is commonly found at relatively higher rate in immunocompromised persons. Cryptosporidiosis in children and immune-compromised individuals such as HIV-infected persons have been reported from many parts of the world. In Nigeria, this too have been reported in children, malnourished and HIV infected individuals, but there is limited data in patients with pTB and T2DM.^{6,7,8}

This study reported very high prevalence of cryptosporidiosis of 26.3% and 13.7%, according to ELISA and mZN technique, respectively. This prevalence is significantly higher than previously estimated

Table I: Prevalence of Cryptosporidiosis among various group of Patients.

Study group	No. of Subjected tested	No. (%) Positive by mZN stain Microscopy	No. Positive (%) by Cryptosporidium antigen ELISA	χ^2	p value
Children (< 5 years)	40	9 (22.5)	25 (62.5)		
HIV infected persons	40	6 (15.0)	8 (20.0)		
Persons with active pTB	40	1 (2.5)	3 (7.5)		
Persons with T2DM	40	6 (15.0)	11 (27.5)		
Total	160	22 (13.8)	42 (26.3)	32.15	<0.0001

mZN - modified Zeihl Neelsen; pTB - Pulmonary Tuberculosis; T2DM—Type 2 Diabetes Mellitus.

Table II: Prevalence of Cryptosporidiosis by Sociodemographic variables of patients presenting with acute diarrhea (n =160).

Sociodemographic characteristic		Negative cases	Positive cases	OR (95%CI)	p value
Age range (years)	<5	15	25	1.67 (0.21-13.01)	0.627
	5 – 14	13	2	0.13 (0.01- 1.80)	0.136
	15 – 24	43	6	0.14 (0.16- 1.18)	0.071
	25 – 34	22	4	0.18 (0.02 -1.69)	0.134
	35 – 44	11	3	0.27 (0.03-2.82)	0.276
	45 – 54	12	0	0.04 (0.00-1.11)	0.057
	≥55	2	2	Referent	
Gender	Male	45	27	2.92 (1.40-6.07)	0.004*
	Female	73	15	Referent	
Educational Level	No Formal education	43	22	3.58 (0.96-13.3)	0.057
	Primary	33	9	.91 (0.46-7.87)	0.371
	High School	21	8	2.67 (0.62-11.46)	0.187
	Tertiary	21	3	Referent	
Residential Area	Rural	25	21	2.62 (0.98-7.03)	0.053
	Suburb	68	13	0.59 (0.22-1.69)	0.309
	Urban	25	8	Referent	
Socioeconomic status	Low	44	29	8.3 (2.36-29.59)	0.0001*
	Middle	38	8	2.67 (0.66-10.8)	0.170
	High	38	3	Referent	
Occupation	Farmer	30	16	0.84 (0.33-2.17)	0.726
	Civil servant	26	3	0.18 (0.05-0.74)	0.017*
	Self-employed	24	6	0.39 (0.13-1.25)	0.114
	Student	19	3	0.25 (0.06-1.03)	0.055
	Unemployed	19	12	Referent	
Marital status	Married	45	27	2.36 (1.10 -5.06)	0.027*
	Widowed	11	1	0.40 (0.05-3.38)	0.402
	Single	62	14	Referent	

Table III: Risk factors of cryptosporidiosis in patients presenting with acute diarrhea (n =160).

Risk Factors		Negative Cases	Positive Cases	OR (95% CI)	p value
Possession of house pet	Yes	23	29	9.21 (4.15-20.4)	<0.0001
	No	95	13	Referent	
Daily consumption vegetables food	Yes	40	31	4.44 (2.02-9.63)	0.0001
	No	72	11	Referent	
Source of Drinking water	Borehole /well	62	23	1.09 (0.54-2.21)	0.804
	Tap	56	19	Referent	
Means of defecation	Open bush	46	22	1.72 (0.85-3.50)	0.133
	Water closet/ pit latrine	72	20	Referent	
Household population size	Dense (≥8 persons)	62	31	2.49 (1.14-5.43)	0.024
	Sparse (<8 persons)	54	11	Referent	
Presence of Household Poultry	Yes	7	15	8.81 (3.27-23.73)	<0.0001
	No	111	27	Referent	

This prevalence is comparable to previous studies reported from Canada (18.02%) and USA (21.2%).^{11,12} Several factors such as age, sex, gastrointestinal conditions, personal and environmental hygiene, sewage water management, consumption of untreated water, contact with infected or carrier animals and poor economic status have been recognized predisposing factors to high prevalence of cryptosporidiosis.

In particular, our data showed that the prevalence was highest among patients aged ≤ 5 years, followed by elderly adults aged ≥ 55 years. This is similar to findings from Khan *et al.*¹³ This age group represent the vulnerable age which usually have deficient immunity due to immature immune function in younger children or wane immunity due to old age.

Out of the 40 diabetic mellitus (DM) patients screened, 27.5% had cryptosporidiosis by ELISA, this finding is similar to the study of Baqai *et al* who reported 25% cryptosporidiosis in DM patients.¹⁴ But lower than 10.6% and 12.9% reported by Cengiz *et al* and Ali *et al*, respectively.^{15,16} The difference could be due to use of less sensitive method in detecting cryptosporidiosis. Indeed, DM is an immunosuppressive disease that could predispose affected person to cryptosporidiosis.

Until recently, cryptosporidiosis was thought to be a rare complication of intestinal disease in people with tuberculosis. Limited case reports and case series documented cryptosporidiosis in person infected with active *Mycobacterium tuberculosis* (MTB). In this study, we found cryptosporidiosis and MTB coinfection in 7.5% patients. This is in consonance to the report by Mor *et al.*¹⁷ We found a significant association between cryptosporidiosis and the study groups in this study, which has not to our knowledge been previously reported. Given overlap in risk fac-

tors for active TB and cryptosporidiosis such as malnutrition, or other immunodeficiency conditions. It is possible that this association is confounded by these or other factors. An association with active TB would support the view that respiratory tract cryptosporidiosis is related to immunocompromised state.

The ELISA positive detectable rate was twice that of the mZN protocol. Similar findings were reported by Cengiz *et al.*¹⁵ It may be impossible to detect the parasite in stool samples, which contain few or distorted oocysts, leading a false negativity of the mZN microscopy. The false negativity of mZN is also due to scarcity of parasite. ELISA kits, however, employing *Cryptosporidium* specific monoclonal antibodies, have a sensitivity and specificity of 93% - 100% and has better chances of detecting the antigen in the stool. The results of cryptosporidium-antigen ELISA indicate that the simple, rapid, reliable, and standardized immunoassay test is sensitive and specific for routine diagnosis and may be useful for large-scale epidemiological studies of cryptosporidiosis.

The high burden of cryptosporidiosis from this study may be attributed to poor personal hygiene and inadequate sanitary facilities. The patients were predominantly illiterates (33.8%) with no formal education, with majority being farmers (34.8%) as occupations, and of low socioeconomic status. For instance, 32.4% of the patients who had no access to a household lavatory and practiced open bush defecation had cryptosporidiosis.

Not using sanitary facilities, water treatment and poor hygiene habits of some of the patients might increase the general environmental contamination and thereby increases the risk of cryptosporidium infection for all other persons living in the same setting.¹⁸ Therefore, the protective effects of sanitation and hygiene habits could have been underestimated. Moreover, owing to its low latitude

and high elevation, the town studied has a mild climate and belongs to the humid sub-tropical zone.

The effects of seasonality in both temperature and rainfall were once shown to be associated with an increased risk of cryptosporidiosis.¹⁸ These local climatic conditions define a typical habitat area for *Cryptosporidium spp.* Alternatively, these differences may reflect the use of different diagnostic techniques. In addition, we found that patients who raised livestock or poultry were more likely to be infected with *Cryptosporidium*, which was consistent with previous study conducted by Yang *et al.*¹⁹

Findings from this study showed that cryptosporidiosis occurred more in male than female. This could be due to the fact that males were exposed to lot of community activities, farming, swimming in the river than the female counterparts in Nigeria. This study is in agreement with a study conducted by Kimani *et al* which showed a wide difference in gender distribution, of males and females.¹⁹ However Tombang *et al* reported a higher prevalence of positive cryptosporidiosis in females than their male counterparts (females, 5.4% versus males, 3.6%), which is in contradiction to both our study and Kimani *et al* findings.²⁰ According to their study, the reason was attributed to the fact that persons of both sexes engage in almost the same recreational activities and so are likely to be equally exposed to similar environmental conditions.

In this study, the prevalence of cryptosporidium infection was higher in HIV infected individuals, when compared to other study in eastern and western Nigeria, this could be due to difference in laboratory testing protocols utilized in these studies and the low socio-economic status of the people in our study area as compared to other localities in the south-east and south-south regions of

Nigeria.^{8,21}

Findings from this study also revealed that individuals who used surface water, well and borehole (for drinking or bathing purposes) were at higher risk for infection, while those who used tap water for consumption were found lesser risk for cryptosporidiosis. Most of the patients were from villages and rural areas where birds, cats and dogs are commonly wandering freely, which may be a route for subsequent zoonotic spreading of oocysts, contaminating the soil and water with their feces. To buttress this, presence of household poultry was found to be significant risk factor of cryptosporidiosis. The observation made from this study showed insignificant association between the detection of *Cryptosporidium* oocysts and the source of drinking water. Though statistically insignificant, the highest number of oocysts was detected among persons from houses dependent on well, bore hole water. Wells in this study area are mostly shallow and can easily be contaminated with human and animal excreta which could serve as the reservoir of the oocysts. This agrees with the reports that contaminated water represents the major source of cryptosporidiosis for humans.^{13,22}

The findings of our study are in agreement that families with low economic status are at significant risk for cryptosporidiosis in the study area. Statistically, there was no significant association observed among the parent level of education and patients' occupation. Comparable study from USA reported that cryptosporidiosis is the highest risk for people living in families with poor food adequacy.¹¹ The food insufficiency may push the families to access food products from places where food safety and hygiene is poor such as at open street marketplaces.¹³

To date, there is no specific therapy developed against human and animal cryptosporidiosis, however, nitazoxanide has shown

promising results and has since been approved by the U.S. FDA as anti-cryptosporidiosis.²³ However, this drug is not yet become globally available. Despite the use of nitazoxanide, its curative effects on HIV/AIDS and other immunocompromised patients are not significantly effective compared to a placebo in some experimental studies.

The strength of our study lies in the use of the more sensitive and specific Sandwich Para-Tech® ELISA kit from *Medical Chemical Corp, California, USA*, which picked up twice the number of positive cases than the usual mZN testing procedure. Thus the prevalence reported by our study using the ELISA technique may represent the true prevalence rate. There are several limitations to our study. First is the population chosen which is specific and unique to our region and hospital and hence results may not be generalizable to the population in general in Nigeria. Second is the lack of multivariate analysis of the positive variables to remove confounding variables which were found to be significant based on univariate analysis.

CONCLUSION

Base on the 15% national prevalence of cryptosporidiosis from a recent metanalysis, it can be inferred that the present study revealed a significantly high prevalence of cryptosporidiosis among patients presenting with acute diarrhea at Abuja Teaching Hospital. It's recommended that government and healthcare policymakers should consider encourage and implement measures that will improve environmental sanitary and socioeconomic conditions of their citizens. In addition, individuals should minimize risk factors associated with personal hygiene identified in this study.

CONFLICT OF INTEREST

None

REFERENCES

- 1: Pielok L, Nowak S, Kludkowska M, Frackowiak K, Kuszel L, Zmora P, et al. [Massive Cryptosporidium infections and chronic diarrhea in HIV-negative patients](#). *Parasitol Res* 2019;118:1937-42. [Accessed on 2020 April 23].
- 2: Shrivastava AK, Kumar S, Smith WA, Sahu PS. [Revisiting the global problem of Cryptosporidiosis and Recommendations](#). *Trop Parasitol*. 2017;7(1):8-17. [Accessed on 2020 April 23].
- 3: Mohammadpour I, Bozag-Chalati F, Motazedian MA. [Molecular characterization and phylogenetic analysis of microsporidia and Cryptosporidium spp. in patients with multiple bowel biopsies from Fars Province](#). *Iran Ann Parasitol*. 2016;62:321-330. [Accessed on 2020 April 23].
- 4: Odeniran PO, Ademola IS. [Epidemiology of Cryptosporidium infection in different hosts in Nigeria: A meta-analysis](#). *Parasitol Int*. 2019;71:194-206. [Accessed on 2020 April 23].
- 5: Cheesbrough M. [District Laboratory Practice in Tropical Countries. Part I](#). Cambridge University Press. 2003;193-208.
- 6: Anejo-Okopi JA, Okojokwu JO, Ebonyi AO, Ejeliogu EU, Isa SE, Audu O, et al. [Molecular characterization of cryptosporidium in children aged 0- 5 years with diarrhea in Jos, Nigeria](#). *Pan Afr Med J*. 2016;25:253. [Accessed on 2020 April 23].
- 7: Aminu M, Ndaks CM, Ella EE. [Cryptosporidium Infection and Correlation with CD4+ T-cell count among Human Immunodeficiency Virus Seropositive Patients within Kaduna Metropolis, Nigeria](#). *Open Conf Proceedings J*. 2014;5:19-28. [Accessed on 2020 April 23].
- 8: Oyedeji OA, Adejuyigbe E, Oninla SA, Akindele AA, Adedokun AA, et al. [Intestinal Parasitoses in HIV Infected Children in a Nigerian Tertiary Hospital](#). *J Clin Diag Res*. 2015;9(11):1-5. [Accessed on 2020 April 23].
- 9: Olopade BO, Ogunniyi TA, Oyekunle AA, Odetoyin BW, Adegoke AO. [Cryptosporidiosis: Prevalence, risk factors and diagnosis in adult HIV-infected patients at Obafemi Awolowo University Teaching Hospitals Complex \(OAUTHC\), Ile-Ife, Osun State, Nigeria](#). *Int J Med Biomed Res* 2017;6(1):18-29. [Accessed on 2020 April 23].
- 10: Saulawa MA, Magaji AA, Faleke O. O, Musawa

- AI, Bala A. Prevalence of Cryptosporidium Coproantigens in Humans in Sokoto State, North Western Nigeria. *Ann Exp Biol.* 2016;4 (2):1-8. [Accessed on 2020 April 23].
 - 11: Becker DJ, Oloya J, Ezeamama AE. [Household socioeconomic and demographic correlates of Cryptosporidium seropositivity in the United States.](#) *PLOS Negl Trop Dis.* 2015;9:e0004080. [Accessed on 2020 April 23].
 - 12: Garcia-R JC, French N, Pita A, Velathanthir N, Shrestha R, Hayman D. [Local and global genetic diversity of protozoan parasites: Spatial distribution of Cryptosporidium and Giardia genotypes.](#) *Plos Negl Trop Dis.* 2017;11:e0005736. [Accessed on 2020 April 23].
 - 13: Khan A, Shams S, Khan S, Khan MI, Khan S, Ali A. [Evaluation of prevalence and risk factors associated with Cryptosporidium infection in rural population of district Buner, Pakistan.](#) *PLoS One.* 2019;14:e0209188. [Accessed on 2020 April 23].
 - 14: Baqai R, Anwar S, Kazmi SU. Detection of cryptosporidium in immunosuppressed patients. *J Ayub Med Coll Abbottabad.* 2005;17 (3):38–40.
 - 15: Cengiz ZT, Hasan Y, Sahin OH, Kapmaz M, Ekici P. [The Frequency of Cryptosporidium spp. in Immunocompromised Patients by Modified Acid-Fast Staining, Cassette Kit and ELISA Methods: Comparison of the Diagnostic Techniques.](#) *Jundishapur J Microbiol.* 2017;10:e36479. [Accessed on 2020 April 23].
 - 16: Ali OS, Mohammad SA, Salman YJ. [Incidence of Some Intestinal Parasites among Diabetic Patients Suffering from Gastroenteritis.](#) *Int J Curr Microbiol App Sci.* 2018;7:3695-3708. [Accessed on 2020 April 23].
 - 17: Mor S, Demaria A, Griffiths J, Naumova EN. [Cryptosporidiosis in the Elderly Population of the United states.](#) *Clin Inf Dis.* 2009;48:698-705. [Accessed on 2020 April 23].
 - 18: Speich B, Croll D, Furst T, Utzinger J, Keiser J. Effect of sanitation and water treatment on intestinal protozoa infection: a systematic review and meta-analysis. *Lancet Infect Dis.* 2016;16:87–99. [Accessed on 2020 April 23].
 - 19: Yang Y, Zhou Y, Xiao PL Shi Y, Chen Y, Liang S, et al. [Prevalence of and risk factors associated with Cryptosporidium infection in an under-developed rural community of southwest China.](#) *Infect Dis Poverty.* 2017;6:2. [Accessed on 2020 April 23].
 - 20: Tombang AN, Ambe NF, Bobga TP, Nkfusai CN, Collins NM, Ngwa SB, et al. [Prevalence and risk factors associated with cryptosporidiosis among children within the ages 0-5 years attending the Limbe regional hospital southwest region, Cameroon.](#) *BMC Public Health.* 2019;19:1144. [Accessed on 2020 April 23].
 - 21: Akinbo FO, Okaka CE, Omoregie R, Dearen T, Leon ET, Xiao L. [Molecular Characterization of Cryptosporidium spp. in HIV-infected Persons in Benin City, Edo State, Nigeria.](#) *Fooyin J Health Sci.* 2010;2:85-89. [Accessed on 2020 April 23].
 - 22: Lv S, Tian L, Liu Q, Qian M, Fu Q, Steinmann P, et al. [Water-related parasitic diseases in China.](#) *Int J Env Res Pub Health.* 2013;10:1977–2016. [Accessed on 2020 April 23].
 - 23: Lee S, Harwood M, Girouard D, Meyers MJ, Campbell MA, Beamer G, et al. [The therapeutic efficacy of azithromycin and nitazoxanide in the acute pig model of Cryptosporidium hominis.](#) *PLOS One.* 2017;12:e0185906. [Accessed on 2020 April 23].
-

APPENDIX I

Definition of Indices of Socioeconomic Status (SES) of Patients Participating in Study.

Persons who had tertiary education, earn more than 200 US\$ per month, resides in urban settlement and works in civil services were considered upper SES, while those that had had tertiary education, earn between 100 - 200 US\$ per month, resides in rural settlement and employed through business were considered middle SES, and persons who had at most high school education, earn less than 100 US\$ per month, resides in rural settlement and works as peasant farmer or unemployed were considered lower SES.

APPENDIX II

Modified Zeihl Neelsen Stain Microscopy:

Detection of *Cryptosporidium* oocysts in the concentrated stool was done using the modified cold Ziehl Neelsen staining technique, as previously described.¹¹ Briefly, a concentrated smear of the stool was made on a clean grease-free slide and fixed in methanol for 3 min. The slide was immersed in cold Carbol fuchsin and stained for 15 min. It was then thoroughly rinsed in tap water and decolorized in 1% HCl (v/v) in methanol for 10 to 15 min. After rinsing again in tap water, the slide was counterstained with 0.4% malachite green for 30 s. The slide was then air-dried and observed under the compound light microscope using 40× objective lens for the presence of *Cryptosporidium* oocysts. Higher-magnification objectives (100× objective) was used to confirm the presence of *Cryptosporidium* spp., oocysts as small pink to red spherules on pale green background.

APPENDIX III

Determination of Fecal *Cryptosporidia* Antigen by Sandwich ELISA

This was done using Para-Tech® ELISA kit from Medical Chemical Corp, California, USA.

This ELISA is an in vitro immunoassay for the qualitative determination of *Cryptosporidium* species antigen in human feces. The assay uses rabbit anti-*Cryptosporidium* polyclonal antibodies to capture the species antigen from the stool supernatant. A second set of goat anti-*Cryptosporidium* polyclonal antibodies were then added which sandwiches the captured species antigen. This reaction was visualized by the addition of anti-goat polyclonal antibodies conjugated to peroxidase and the chromogen tetramethylbenzidine (TMB). The resulting blue color development was converted to an easily read yellow color by addition of an acidic "stop solution" to end the reaction. The presence of yellow color above 0.15 Optical Density of absorbance indicates presence of *Cryptosporidium* oocysts or species antigens. This procedure has Sensitivity of 100% (95% CI: 86% - 100%) and Specificity of 97% (95% CI: 93% to 99%). *Cryptosporidiosis* was defined by positive ELISA result.